

## **Introduction**

Section 4121.44 of the Ohio Revised Code requires the Ohio Bureau of Workers' Compensation (BWC) to collect health care data for the purpose of measuring and analyzing "costs, quality, appropriateness of medical care, and effectiveness of medical care delivered by all components of the workers' compensation system."

Part of these measures include a "provider outcome measurement" for evaluating health care providers, according to Ohio Administrative Code section 4123-6-02 (T), on the basis of cost, utilization, and treatment outcomes efficiency and compliance with bureau requirements.

This is achieved by analyzing (at a minimum) line item detail from a medical bill and employee specific information including, but not limited to:

- Demographics;
- Diagnosis allowances;
- Return-to-work and remain-at-work statistics;
- Other data regarding treatment.

## **Description of results**

To perform this required data analysis and to create benchmarks for establishing provider outcome measurements, BWC contracted with the Work Loss Data Institute to analyze BWC claims with dates of injury between Jan. 1, 2000 and Dec. 31, 2003.

Claims in the analysis include roughly 925,000 claims managed by managed care programs (MCOs) through BWC's Health Partnership Program. Excluded were claims managed by self-insured employers, bankrupt self-insured employers, or black lung and marine fund programs.

The attached spreadsheets represent the disability duration outcomes received from Work Loss Data Institute.

The first spreadsheet, titled "Ohio Specific Workers' Compensation Disability Durations per Most Severe ICD-9", provides outcomes for ICD-9s which were listed as the "most severe" in at least 15 claims.

The second spreadsheet, titled "Ohio Workers' Compensation Specific Disability Durations per Common Select Surgical Procedures," reports the duration of disability outcomes following performance of the given CPT codes until the injured worker returns to work."

(Due to licensing restrictions, BWC cannot provide the description of the CPT Code.)

When reviewing these outcomes, it is important to understand the concepts and limitations of the results. Important factors include the following items:

1. The outcomes represent BWC-specific results for the claims allowed during the study period and treated within the rules and processes of the Ohio workers' compensation system. **The results represent actual measurements and not consensus durations of disability, practicing under optimal conditions, or private pay insurance results which may be found in other benchmarks.** While benchmarks are provided such as the Official Disability Guidelines (ODG) and OSHA, those benchmarks do not represent Ohio-specific and perhaps not workers' compensation benchmarks.
2. The Ohio workers' compensation system is based on conditions (ICD-9 diagnoses) specifically recognized in the claim. In this analysis, all lost work days are assigned to the ICD-9 code determined to be "most severe" based on a two-pass process. Initially, all lost days were assigned to each ICD-9 code allowed in the claim. From the entire population of claims, this pass created a distribution of the actual number of lost work days for each ICD-9 in the population of all claims. This distribution was used to determine the median disability duration for each ICD-9 code. A second pass through the population of claims assigned the number of lost work days in the claim to the ICD-9 code with the greatest median disability duration allowed in the claim.
3. The spreadsheet for Ohio Specific Workers' Compensation Disability Durations per Most Severe ICD-9 includes the total number of claims in which the ICD-9 code was considered most severe. It then provides the number and percentage of claims in which:
  - There were zero lost work days;
  - The injured worker has never returned to work or no return to work data was available;
  - There were one or more lost work days.

It also provides the number of days required to return a given decile of those individuals who returned to work but who had missed one or more days back to work. The Ohio workers' compensation median and mean are provided, along with the ODG and OSHA medians when available.

A. It is important to realize that the return-to-work data includes only those claims which had one or more lost work days. This population also includes claims for those individuals undergoing surgery, so lost work days following surgery are included.

B. Claims were assigned to the most severe ICD-9 code at the three-, four- and five-digit level. BWC typically codes only to the more specific five-digit level where one exists. Therefore, many codes indicated in the outcome results may not be recognized by BWC and are indicated in the third column by an "I" ("Invalid" for BWC). Claims in these "invalid" codes also are found in the "valid" five-digit ICD-9 codes. For example, a claim in which the most severe ICD-9 is a herniated lumbar disc (ICD-9 code 722.10) will be found in ICDs 722, 722.1 and 722.10. Only 722.10 is considered valid by BWC.

4. The spreadsheet for Ohio Workers' Compensation Specific Disability Durations per Common Select Surgical Procedures is limited to common select procedures used to treat musculoskeletal conditions on an elective basis. For each specific CPT code, the numbers of procedures are indicated; along with the number of days required from the date of the procedure for a given decile of individuals undergoing the procedure to return to work. A given procedure may be used more than once in a claim or multiple procedures may be found in the same claim. The number and percent of "open cases" represent those claims in which the injured worker has not returned to work, or for which no return-to-work date is available in the claim file.
5. Many confounding factors are not available in the claims data, or could not be included in the analysis. An earlier analysis of the data found poor correlation between the duration of disability and the level of work activity performed by the individual at the time of injury, to the extent that the level of work activity was not considered in this particular analysis. On an individual case basis, it is believed that the level of work activity to which the individual must return is an important return-to-work factor. However, on this population-based outcome analysis, this assumption could not be supported. Among the reasons this assumption may not have been validated in this population include:
  - The fact that many injured workers may have returned to less physically demanding jobs;
  - The specific level of work activity may have been misclassified;
  - Motivation, other confounding factors and psychosocial factors could not be addressed by the analyses.

## Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

Notes: \* Data represents Ohio state fund claims with dates of injury from Jan. 1, 2000 through Dec. 31, 2003.

\* Data includes claims treated surgically.

\* Claims assigned to ICD-9 considered "most severe" based on statistical analysis.

\* ICD-9 codes with fewer than 15 cases not included in analysis.

\* ICD-9 Codes considered "valid" for reimbursement/allowance by BWC indicated in **bold**.

\* "Invalid ICD-9" indicates ICD-9 Code usually of three or four digits that are not recognized by BWC, since BWC usually codes to five-digit ICD-9. These "invalid codes" shown by normal font.

\* Number of claims include claims with zero lost days, with one or more last days, and those in which there has been no return to work.

\* Disability duration percentile determined using cases with one or more days lost work time in which there has been a return to work.

\* ICD-9 Codes found in fewer than 200 claims may be less precise, due to the relative fewer number of claims in the sample size. ICD-9 codes with more than 200 claims in **bold**.

Disability Durations by ICD9 Diagnosis Code, 0-Day Waiting Period, Including Surgeries																							
ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
041	I	Bacterial infection in conditions classified elsewhere and of unspecified site	28	2	7%	24	86%	15	20	24	27	30	35	46	138	176	287	30	66.38	8		2	7%
041.1	I	Staphylococcus	17	1	6%	15	88%	15	20	24	27	32	35	83	138	176	239	32	69.07	14		1	6%
110	I	Dermatophytosis	69	49	71%	15	22%	1	1	2	2	4	4	6	6	16	19	4	5.47	2		5	7%
133	I	Acariasis	196	123	63%	62	32%	1	1	1	2	2	3	4	6	9	32	2	4.48	3	2	11	6%
<b>133.0</b>		<b>Scabies</b>	<b>189</b>	<b>118</b>	<b>62%</b>	<b>60</b>	<b>32%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>9</b>	<b>32</b>	<b>2</b>	<b>4.57</b>	<b>3</b>		<b>11</b>	<b>6%</b>
215	I	Other benign neoplasm of connective and other soft tissue	45	9	20%	27	60%	6	22	60	81	165	211	320	365	365	365	165	172.37	14		9	20%
<b>215.2</b>		<b>Upper limb, including shoulder</b>	<b>27</b>	<b>4</b>	<b>15%</b>	<b>16</b>	<b>59%</b>	<b>6</b>	<b>23</b>	<b>60</b>	<b>81</b>	<b>173</b>	<b>326</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>173</b>	<b>196.00</b>	<b>11</b>		<b>7</b>	<b>26%</b>
276	I	Disorders of fluid, electrolyte, and acid-base balance	87	48	55%	35	40%	1	1	1	1	2	2	4	5	9	37	2	4.66	8		4	5%
276.5	I	Volume depletion	83	46	55%	34	41%	1	1	1	1	2	3	4	6	9	37	2	4.74	6		3	4%
296	I	Episodic mood disorders	926	14	2%	161	17%	20	85	145	203	292	365	365	365	365	365	292	238.62	10		751	81%
296.2	I	Major depressive disorder, single episode	760	8	1%	135	18%	18	80	135	187	292	365	365	365	365	365	292	236.40	21		617	81%
<b>296.20</b>		<b>Major depressive disorder, single episode</b>	<b>183</b>	<b>2</b>	<b>1%</b>	<b>44</b>	<b>24%</b>	<b>9</b>	<b>34</b>	<b>102</b>	<b>167</b>	<b>203</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>203</b>	<b>218.02</b>	<b>21</b>		<b>137</b>	<b>75%</b>
<b>296.22</b>		<b>Major depressive disorder, single episode</b>	<b>244</b>	<b>2</b>	<b>1%</b>	<b>46</b>	<b>19%</b>	<b>18</b>	<b>74</b>	<b>122</b>	<b>160</b>	<b>258</b>	<b>334</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>258</b>	<b>224.24</b>	<b>21</b>		<b>196</b>	<b>80%</b>
<b>296.23</b>		<b>Major depressive disorder, single episode</b>	<b>292</b>	<b>3</b>	<b>1%</b>	<b>38</b>	<b>13%</b>	<b>55</b>	<b>112</b>	<b>192</b>	<b>275</b>	<b>292</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>292</b>	<b>258.66</b>	<b>21</b>		<b>251</b>	<b>86%</b>
296.3	I	Major depressive disorder, recurrent episode	128	1	1%	23	18%	43	102	165	253	365	365	365	365	365	365	365	259.91	18		104	81%
300	I	Anxiety, dissociative and somatoform disorders	344	10	3%	99	29%	6	25	51	136	177	233	313	365	365	365	177	185.35	11	47	235	68%
300.0	I	Anxiety states	117	4	3%	36	31%	3	17	35	78	136	185	313	365	365	365	136	164.72	11		77	66%
<b>300.00</b>		<b>Anxiety state, unspecified</b>	<b>49</b>	<b>2</b>	<b>4%</b>	<b>16</b>	<b>33%</b>	<b>10</b>	<b>38</b>	<b>41</b>	<b>90</b>	<b>136</b>	<b>151</b>	<b>313</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>136</b>	<b>169.69</b>	<b>41</b>	<b>24</b>	<b>31</b>	<b>63%</b>
<b>300.4</b>		<b>Dysthymic disorder</b>	<b>197</b>	<b>5</b>	<b>3%</b>	<b>57</b>	<b>29%</b>	<b>7</b>	<b>29</b>	<b>90</b>	<b>149</b>	<b>179</b>	<b>246</b>	<b>299</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>179</b>	<b>192.72</b>	<b>26</b>		<b>135</b>	<b>69%</b>
307	I	Special symptoms or syndromes, not elsewhere classified	168	6	4%	35	21%	12	28	54	84	109	190	231	303	365	365	109	159.26	13		127	76%
307.8	I	Pain disorders related to psychological factors	166	6	4%	34	20%	17	28	69	86	109	206	231	365	365	365	109	163.82	3		126	76%
<b>307.89</b>		<b>Other</b>	<b>154</b>	<b>2</b>	<b>1%</b>	<b>28</b>	<b>18%</b>	<b>19</b>	<b>69</b>	<b>86</b>	<b>109</b>	<b>145</b>	<b>209</b>	<b>237</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>145</b>	<b>185.21</b>	<b>19</b>		<b>124</b>	<b>81%</b>
308	I	Acute reaction to stress	117	8	7%	71	61%	14	26	44	57	71	103	147	229	365	365	71	125.85	9	35	38	32%
<b>308.3</b>		<b>Other acute reactions to stress</b>	<b>105</b>	<b>7</b>	<b>7%</b>	<b>61</b>	<b>58%</b>	<b>14</b>	<b>28</b>	<b>44</b>	<b>57</b>	<b>70</b>	<b>100</b>	<b>147</b>	<b>229</b>	<b>365</b>	<b>365</b>	<b>70</b>	<b>122.80</b>	<b>9</b>		<b>37</b>	<b>35%</b>
309	I	Adjustment reaction	730	25	3%	303	42%	18	47	72	97	136	193	275	365	365	365	136	175.81	13	4	402	55%
<b>309.0</b>		<b>Adjustment disorder with depressed mood</b>	<b>129</b>	<b>3</b>	<b>2%</b>	<b>47</b>	<b>36%</b>	<b>39</b>	<b>86</b>	<b>128</b>	<b>165</b>	<b>195</b>	<b>318</b>	<b>359</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>195</b>	<b>220.94</b>	<b>51</b>		<b>79</b>	<b>61%</b>
309.2	I	With predominant disturbance of other emotions	241	11	5%	84	35%	7	42	74	106	139	223	339	365	365	365	139	185.71	12		146	61%
<b>309.28</b>		<b>Adjustment disorder with mixed anxiety and depressed mood</b>	<b>214</b>	<b>8</b>	<b>4%</b>	<b>70</b>	<b>33%</b>	<b>9</b>	<b>53</b>	<b>74</b>	<b>106</b>	<b>167</b>	<b>240</b>	<b>342</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>167</b>	<b>193.64</b>	<b>68</b>		<b>136</b>	<b>64%</b>
309.8	I	Other specified adjustment reactions	321	11	3%	156	49%	18	38	60	83	107	146	208	328	365	365	107	153.90	14		154	48%
<b>309.81</b>		<b>Posttraumatic stress disorder</b>	<b>311</b>	<b>11</b>	<b>4%</b>	<b>148</b>	<b>48%</b>	<b>15</b>	<b>38</b>	<b>61</b>	<b>83</b>	<b>110</b>	<b>167</b>	<b>209</b>	<b>333</b>	<b>365</b>	<b>365</b>	<b>110</b>	<b>155.34</b>	<b>35</b>		<b>152</b>	<b>49%</b>
310	I	Specific nonpsychotic mental disorders due to brain damage	372	95	26%	224	60%	2	3	5	9	13	19	36	62	134	365	13	47.22	16		53	14%
<b>310.2</b>		<b>Postconcussion syndrome</b>	<b>355</b>	<b>95</b>	<b>27%</b>	<b>217</b>	<b>61%</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>12</b>	<b>17</b>	<b>29</b>	<b>54</b>	<b>121</b>	<b>365</b>	<b>12</b>	<b>42.05</b>	<b>22</b>		<b>43</b>	<b>12%</b>
<b>311</b>		<b>Depressive disorder, not elsewhere classified</b>	<b>801</b>	<b>24</b>	<b>3%</b>	<b>165</b>	<b>21%</b>	<b>42</b>	<b>94</b>	<b>149</b>	<b>209</b>	<b>278</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>278</b>	<b>241.95</b>	<b>21</b>		<b>612</b>	<b>76%</b>
337	I	Disorders of the autonomic nervous system	468	15	3%	198	42%	22	67	124	183	251	365	365	365	365	365	251	229.73	22		255	54%
337.2	I	Reflex sympathetic dystrophy	460	14	3%	194	42%	24	67	130	183	251	365	365	365	365	365	251	230.24	109		252	55%
<b>337.21</b>		<b>Reflex sympathetic dystrophy of the upper limb</b>	<b>246</b>	<b>7</b>	<b>3%</b>	<b>110</b>	<b>45%</b>	<b>24</b>	<b>91</b>	<b>135</b>	<b>179</b>	<b>296</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>296</b>	<b>236.82</b>	<b>109</b>		<b>129</b>	<b>52%</b>

### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open	
337.22		<b>Reflex sympathetic dystrophy of the lower limb</b>	199	5	3%	73	37%	41	81	135	201	254	365	365	365	365	365	254	237.62	111		121	61%	
353	I	Nerve root and plexus disorders	216	54	25%	103	48%	5	19	45	84	99	157	257	365	365	365	99	158.47	12		59	27%	
353.0		<b>Brachial plexus lesions</b>	120	29	24%	59	49%	6	31	52	90	108	164	257	365	365	365	108	162.93	38		32	27%	
353.8		<b>Other nerve root and plexus disorders</b>	73	17	23%	39	53%	2	13	44	52	92	137	332	365	365	365	92	160.62	44		17	23%	
354	I	Mononeuritis of upper limb and mononeuritis multiplex	6,769	1,697	25%	4,156	61%	14	29	42	53	67	84	110	159	278	365	67	103.26	13		916	14%	
354.0		<b>Carpal tunnel syndrome</b>	5,601	1,432	26%	3,514	63%	15	29	41	52	64	81	100	139	229	365	64	95.30	24	32	655	12%	
354.1		<b>Other lesion of median nerve</b>	104	35	34%	53	51%	3	10	31	54	69	103	165	222	365	365	69	119.00	10		16	15%	
354.2		<b>Lesion of ulnar nerve</b>	943	203	22%	524	56%	11	31	49	68	96	138	214	323	365	365	96	148.59	43		216	23%	
354.3		<b>Lesion of radial nerve</b>	102	23	23%	53	52%	10	25	49	102	120	181	273	360	365	365	120	165.11	42		26	25%	
355	I	Mononeuritis of lower limb	293	51	17%	162	55%	5	27	45	68	102	153	249	365	365	365	102	155.03	22		80	27%	
355.3		<b>Lesion of lateral popliteal nerve</b>	33	5	15%	18	55%	1	99	153	184	249	260	320	365	365	365	249	216.61	104		10	30%	
355.5		<b>Tarsal tunnel syndrome</b>	60	5	8%	32	53%	19	40	61	66	92	138	305	365	365	365	92	163.78	61		23	38%	
355.6		<b>Lesion of plantar nerve</b>	40	8	20%	23	58%	19	40	75	92	97	120	213	314	334	365	97	145.87	31		9	23%	
355.7	I	Other mononeuritis of lower limb	27		0%	18	67%	12	27	42	45	46	97	171	325	365	365	46	132.11	22		9	33%	
355.79		<b>Other mononeuritis of lower limb</b>	22		0%	15	68%	12	13	34	42	45	46	150	214	365	365	45	116.33	34		7	32%	
355.8		<b>Mononeuritis of lower limb, unspecified</b>	73	15	21%	40	55%	1	14	31	46	67	112	202	365	365	365	67	144.15	42		18	25%	
355.9		<b>Mononeuritis of unspecified site</b>	31	11	35%	16	52%	4	19	32	97	109	118	230	269	365	365	109	145.75	22		4	13%	
360	I	Disorders of the globe	68	46	68%	22	32%	1	1	2	2	2	5	7	8	34	250	2	17.91	5			0%	
361	I	Retinal detachments and defects	39	9	23%	26	67%	3	11	19	25	37	58	88	117	154	282	37	63.85	22		4	10%	
362	I	Other retinal disorders	27	10	37%	16	59%	2	3	4	7	12	14	20	26	50	158	12	22.81	15		1	4%	
364	I	Disorders of iris and ciliary body	425	230	54%	179	42%	1	1	2	3	4	5	7	11	17	365	4	12.42	5		16	4%	
364.0	I	Acute and subacute iridocyclitis	217	132	61%	77	35%	1	1	2	3	4	4	5	9	12	146	4	7.48	5		8	4%	
364.00		<b>Acute and subacute iridocyclitis, unspecified</b>	147	88	60%	53	36%	1	2	2	3	4	4	5	7	10	146	4	8.49	4		6	4%	
364.3		<b>Unspecified iridocyclitis</b>	134	81	60%	50	37%	1	1	2	2	3	4	4	5	7	12	31	3	5.60	5		3	2%
364.4	I	Vascular disorders of iris and ciliary body	55	11	20%	42	76%	2	4	5	5	6	8	11	13	16	31	6	8.55	11		2	4%	
364.41		<b>Hyphema</b>	54	11	20%	41	76%	2	4	5	6	6	8	11	13	16	31	6	8.63	11		2	4%	
366	I	Cataract	55	10	18%	36	65%	7	12	17	44	55	74	129	159	365	365	55	102.06	14		9	16%	
366.2	I	Traumatic cataract	45	9	20%	30	67%	4	11	13	38	46	58	106	129	165	365	46	82.30	21		6	13%	
366.20		<b>Traumatic cataract, unspecified</b>	27	7	26%	18	67%	7	12	13	38	48	58	74	129	133	165	48	59.94	55		2	7%	
368	I	Visual disturbances	84	42	50%	35	42%	1	1	2	9	14	19	40	60	197	365	14	56.14	7	3	7	8%	
369	I	Blindness and low vision	68	8	12%	44	65%	2	20	58	77	97	129	180	243	351	365	97	133.27	13	180	16	24%	
369.6	I	Profound impairment, one eye	22		0%	15	68%	1	20	76	81	97	105	129	180	243	365	97	117.53	13		7	32%	
370	I	Keratitis	3,062	1,774	58%	1,171	38%	1	1	1	1	2	2	3	4	5	364	2	3.14	2		117	4%	
370.0	I	Corneal ulcer	173	110	64%	56	32%	1	1	2	2	3	4	5	6	12	45	3	5.39	5		7	4%	
370.00		<b>Corneal ulcer, unspecified</b>	129	84	65%	42	33%	1	1	1	2	3	4	5	9	13	45	3	6.17	3		3	2%	
370.2	I	Superficial keratitis without conjunctivitis	2,138	1,210	57%	838	39%	1	1	1	1	2	2	3	3	4	364	2	2.83	2		90	4%	
370.20		<b>Superficial keratitis, unspecified</b>	179	125	70%	46	26%	1	1	1	2	2	3	4	6	33	2	3.54	3		8	4%		
370.21		<b>Punctate keratitis</b>	149	98	66%	43	29%	1	1	1	2	2	2	3	4	14	2	2.47	2		8	5%		
370.24		<b>Photokeratitis</b>	1,806	985	55%	747	41%	1	1	1	1	2	2	3	3	4	364	2	2.80	2	1	74	4%	
370.4	I	Other and unspecified keratoconjunctivitis	147	95	65%	49	33%	1	1	1	2	2	3	4	6	11	20	2	3.76	2		3	2%	
370.40		<b>Keratoconjunctivitis, unspecified</b>	116	76	66%	37	32%	1	1	1	2	2	3	3	6	11	20	2	3.86	3		3	3%	
370.8		<b>Other forms of keratitis</b>	111	64	58%	46	41%	1	1	1	2	2	2	3	3	5	12	2	2.39	2		1	1%	
370.9		<b>Unspecified keratitis</b>	446	264	59%	167	37%	1	1	1	2	2	3	3	4	5	219	2	4.07	2		15	3%	
371	I	Corneal opacity and other disorders of cornea	196	126	64%	60	31%	1	2	2	3	4	5	10	14	32	263	4	18.37	14		10	5%	
371.0	I	Corneal scars and opacities	46	29	63%	16	35%	1	1	2	3	4	5	5	8	18	18	4	5.69	8		1	2%	
371.2	I	Corneal edema	57	31	54%	20	35%	1	2	2	3	6	10	12	19	34	243	6	25.20	10		6	11%	
371.20		<b>Corneal edema, unspecified</b>	48	27	56%	16	33%	1	2	2	4	6	10	12	13	26	34	6	9.56	6		5	10%	
371.4	I	Corneal degenerations	49	31	63%	17	35%	1	2	2	3	4	4	4	9	34	78	4	10.06	8		1	2%	
371.42		<b>Recurrent erosion of cornea</b>	42	27	64%	15	36%	1	1	2	2	4	4	4	9	34	78	4	10.87	4			0%	
372	I	Disorders of conjunctiva	9,020	7,002	78%	1,747	19%	1	1	1	1	2	2	3	4	5	365	2	4.26	2	1	271	3%	
372.0	I	Acute conjunctivitis	5,135	3,962	77%	1,043	20%	1	1	1	1	2	2	3	4	5	365	2	4.17	2	3	130	3%	

### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
<b>372.00</b>		<b>Acute conjunctivitis, unspecified</b>	<b>935</b>	<b>723</b>	<b>77%</b>	<b>182</b>	<b>19%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>365</b>	<b>2</b>	<b>4.56</b>	<b>2</b>		<b>30</b>	<b>3%</b>
<b>372.01</b>		<b>Serous conjunctivitis, except viral</b>	<b>143</b>	<b>103</b>	<b>72%</b>	<b>36</b>	<b>25%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>4</b>	<b>6</b>	<b>2</b>	<b>1.89</b>	<b>2</b>		<b>4</b>	<b>3%</b>
<b>372.03</b>		<b>Other mucopurulent conjunctivitis</b>	<b>177</b>	<b>147</b>	<b>83%</b>	<b>23</b>	<b>13%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>10</b>	<b>2</b>	<b>2.70</b>	<b>2</b>		<b>7</b>	<b>4%</b>
<b>372.05</b>		<b>Acute atopic conjunctivitis</b>	<b>3,866</b>	<b>2,979</b>	<b>77%</b>	<b>800</b>	<b>21%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>365</b>	<b>2</b>	<b>4.22</b>	<b>2</b>		<b>87</b>	<b>2%</b>
372.1	I	Chronic conjunctivitis	130	101	78%	23	18%	1	1	1	1	2	2	3	9	34	214	2	16.26	3		6	5%
<b>372.14</b>		<b>Other chronic allergic conjunctivitis</b>	<b>119</b>	<b>94</b>	<b>79%</b>	<b>19</b>	<b>16%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>34</b>	<b>214</b>	<b>2</b>	<b>15.00</b>	<b>3</b>		<b>6</b>	<b>5%</b>
372.3	I	Other and unspecified conjunctivitis	3,124	2,452	78%	562	18%	1	1	1	1	2	2	3	4	5	365	2	4.02	2		110	4%
<b>372.30</b>		<b>Conjunctivitis, unspecified</b>	<b>2,520</b>	<b>1,943</b>	<b>77%</b>	<b>484</b>	<b>19%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>365</b>	<b>2</b>	<b>4.28</b>	<b>2</b>		<b>93</b>	<b>4%</b>
<b>372.39</b>		<b>Other</b>	<b>602</b>	<b>507</b>	<b>84%</b>	<b>78</b>	<b>13%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>11</b>	<b>2</b>	<b>2.40</b>	<b>2</b>		<b>17</b>	<b>3%</b>
372.7	I	Conjunctival vascular disorders and cysts	539	422	78%	98	18%	1	1	1	1	2	2	3	3	6	81	2	3.34	2		19	4%
<b>372.72</b>		<b>Conjunctival hemorrhage</b>	<b>432</b>	<b>338</b>	<b>78%</b>	<b>79</b>	<b>18%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>6</b>	<b>81</b>	<b>2</b>	<b>3.52</b>	<b>2</b>		<b>15</b>	<b>3%</b>
<b>372.73</b>		<b>Conjunctival edema</b>	<b>64</b>	<b>48</b>	<b>75%</b>	<b>15</b>	<b>23%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>5</b>	<b>14</b>	<b>2</b>	<b>2.73</b>	<b>2</b>		<b>1</b>	<b>2%</b>
373	I	Inflammation of eyelids	357	267	75%	71	20%	1	1	1	2	2	3	3	4	6	49	2	4.00	3	1	19	5%
373.0	I	Blepharitis	129	102	79%	21	16%	1	1	1	1	2	3	3	4	12	27	3	4.43	3		6	5%
<b>373.00</b>		<b>Blepharitis, unspecified</b>	<b>117</b>	<b>94</b>	<b>80%</b>	<b>19</b>	<b>16%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>13</b>	<b>27</b>	<b>3</b>	<b>4.21</b>	<b>3</b>		<b>4</b>	<b>3%</b>
373.1	I	Hordeolum and other deep inflammation of eyelid	140	98	70%	32	23%	1	1	1	2	2	2	3	4	6	49	2	4.16	2		10	7%
<b>373.11</b>		<b>Hordeolum externum</b>	<b>95</b>	<b>65</b>	<b>68%</b>	<b>21</b>	<b>22%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>11</b>	<b>2</b>	<b>2.29</b>	<b>2</b>		<b>9</b>	<b>9%</b>
374	I	Other disorders of eyelids	77	50	65%	24	31%	2	2	3	4	15	31	48	197	365	365	15	79.75	33		3	4%
379	I	Other disorders of eye	534	376	70%	140	26%	1	1	2	2	3	4	6	25	60	275	3	23.20	4		18	3%
379.0	I	Scleritis and episcleritis	58	39	67%	17	29%	1	1	1	2	2	2	2	4	6	6	2	2.41	4		2	3%
379.2	I	Disorders of vitreous body	74	43	58%	29	39%	4	4	12	20	38	45	56	65	192	256	38	50.79	27		2	3%
379.23	I	Vitreous hemorrhage	28	10	36%	17	61%	4	12	33	38	45	49	56	65	92	256	45	52.76	40		1	4%
379.9	I	Unspecified disorder of eye and adnexa	314	237	75%	74	24%	1	1	1	2	2	2	3	4	6	27	2	3.30	3	2	3	1%
<b>379.91</b>		<b>Pain in or around eye</b>	<b>102</b>	<b>74</b>	<b>73%</b>	<b>28</b>	<b>27%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>8</b>	<b>2</b>	<b>2.57</b>	<b>3</b>			<b>0%</b>
<b>379.93</b>		<b>Redness or discharge of eye</b>	<b>78</b>	<b>60</b>	<b>77%</b>	<b>17</b>	<b>22%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>16</b>	<b>25</b>	<b>2</b>	<b>4.06</b>	<b>3</b>		<b>1</b>	<b>1%</b>
<b>379.99</b>		<b>Other ill-defined disorders of eye</b>	<b>93</b>	<b>73</b>	<b>78%</b>	<b>18</b>	<b>19%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>7</b>	<b>27</b>	<b>1</b>	<b>3.61</b>	<b>3</b>		<b>2</b>	<b>2%</b>
384	I	Other disorders of tympanic membrane	62	43	69%	18	29%	1	1	2	2	2	3	6	17	99	120	2	16.72	4		1	2%
384.2	I	Perforation of tympanic membrane	61	43	70%	17	28%	1	1	2	2	3	6	6	17	99	120	3	17.65	6		1	2%
386	I	Vertiginous syndromes and other disorders of vestibular system	50	8	16%	29	58%	6	18	27	61	109	118	133	180	365	365	109	119.66	4		13	26%
386.1	I	Other and unspecified peripheral vertigo	25	3	12%	17	68%	6	18	27	35	68	109	118	150	244	365	68	94.12	6		5	20%
388	I	Other disorders of ear	252	183	73%	44	17%	1	2	2	3	5	7	9	26	32	199	5	15.32	2		25	10%
388.3	I	Tinnitus	97	73	75%	20	21%	1	1	2	2	3	4	6	9	28	199	3	16.55	5	3	4	4%
<b>388.30</b>		<b>Tinnitus, unspecified</b>	<b>84</b>	<b>62</b>	<b>74%</b>	<b>18</b>	<b>21%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>4</b>	<b>9</b>	<b>20</b>	<b>32</b>	<b>199</b>	<b>2</b>	<b>17.89</b>	<b>4</b>		<b>4</b>	<b>5%</b>
389	I	Hearing loss	195	144	74%	35	18%	2	3	9	17	38	47	55	98	274	365	38	77.34	4	10	16	8%
410	I	Acute myocardial infarction	40	4	10%	22	55%	16	31	55	60	65	106	128	181	365	365	65	118.32	31	21	14	35%
415	I	Acute pulmonary heart disease	57	3	5%	47	82%	30	65	91	103	138	175	268	316	365	365	138	175.28	16		7	12%
415.1	I	Pulmonary embolism and infarction	56	3	5%	46	82%	33	67	91	110	138	175	282	316	365	365	138	178.74	27		7	13%
<b>415.19</b>		<b>Other</b>	<b>47</b>	<b>3</b>	<b>6%</b>	<b>40</b>	<b>85%</b>	<b>33</b>	<b>65</b>	<b>83</b>	<b>100</b>	<b>136</b>	<b>175</b>	<b>268</b>	<b>328</b>	<b>365</b>	<b>365</b>	<b>136</b>	<b>179.75</b>	<b>33</b>		<b>4</b>	<b>9%</b>
444	I	Arterial embolism and thrombosis	32	2	6%	26	81%	28	36	41	47	59	91	102	233	365	365	59	120.62	10		4	13%
444.2	I	Of arteries of the extremities	27	2	7%	21	78%	28	34	41	46	59	68	92	131	315	365	59	105.52	25		4	15%
451	I	Phlebitis and thrombophlebitis	92	15	16%	65	71%	4	8	14	21	38	54	78	126	326	365	38	86.49	17	12	12	13%
<b>451.0</b>		<b>Of superficial vessels of lower extremities</b>	<b>26</b>	<b>8</b>	<b>31%</b>	<b>17</b>	<b>65%</b>	<b>2</b>	<b>4</b>	<b>6</b>	<b>6</b>	<b>20</b>	<b>21</b>	<b>24</b>	<b>28</b>	<b>61</b>	<b>273</b>	<b>20</b>	<b>32.53</b>	<b>27</b>		<b>1</b>	<b>4%</b>
451.1	I	Of deep vessels of lower extremities	42	2	5%	35	83%	10	14	39	54	73	98	126	216	365	365	73	121.11	27		5	12%
<b>451.19</b>		<b>Other</b>	<b>30</b>	<b>1</b>	<b>3%</b>	<b>26</b>	<b>87%</b>	<b>10</b>	<b>27</b>	<b>52</b>	<b>55</b>	<b>73</b>	<b>98</b>	<b>115</b>	<b>193</b>	<b>326</b>	<b>365</b>	<b>73</b>	<b>110.69</b>	<b>34</b>		<b>3</b>	<b>10%</b>
453	I	Other venous embolism and thrombosis	143	11	8%	106	74%	14	23	29	47	65	99	121	202	319	365	65	109.18	12		26	18%
<b>453.8</b>		<b>Of other specified veins</b>	<b>118</b>	<b>9</b>	<b>8%</b>	<b>89</b>	<b>75%</b>	<b>14</b>	<b>23</b>	<b>29</b>	<b>41</b>	<b>65</b>	<b>86</b>	<b>116</b>	<b>170</b>	<b>270</b>	<b>365</b>	<b>65</b>	<b>102.67</b>	<b>37</b>		<b>20</b>	<b>17%</b>
<b>453.9</b>		<b>Of unspecified site</b>	<b>20</b>	<b>2</b>	<b>10%</b>	<b>16</b>	<b>80%</b>	<b>13</b>	<b>25</b>	<b>31</b>	<b>57</b>	<b>99</b>	<b>143</b>	<b>253</b>	<b>281</b>	<b>365</b>	<b>365</b>	<b>99</b>	<b>148.75</b>	<b>11</b>		<b>2</b>	<b>10%</b>
<b>459</b>		<b>Other disorders of circulatory system</b>	<b>29</b>	<b>11</b>	<b>38%</b>	<b>17</b>	<b>59%</b>	<b>2</b>	<b>5</b>	<b>5</b>	<b>9</b>	<b>15</b>	<b>34</b>	<b>43</b>	<b>91</b>	<b>254</b>	<b>365</b>	<b>15</b>	<b>61.71</b>	<b>13</b>		<b>1</b>	<b>3%</b>
459.8	I	Other specified disorders of circulatory system	29	11	38%	17	59%	2	5	5	9	15	34	43	91	254	365	15	61.71	14		1	3%
<b>470</b>		<b>Deviated nasal septum</b>	<b>67</b>	<b>26</b>	<b>39%</b>	<b>40</b>	<b>60%</b>	<b>3</b>	<b>8</b>	<b>9</b>	<b>13</b>	<b>16</b>	<b>19</b>	<b>25</b>	<b>39</b>	<b>44</b>	<b>112</b>	<b>16</b>	<b>23.93</b>	<b>13</b>		<b>1</b>	<b>1%</b>
477	I	Allergic rhinitis	64	44	69%	17	27%	1	2	2	3	5	8	13	30	43	158	5	19.47	3	3	3	5%

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
478	I	Other diseases of upper respiratory tract	37	15	41%	19	51%	1	5	8	11	33	38	101	170	365	365	33	86.74	10	20	3	8%
<b>486</b>		<b>Pneumonia, organism unspecified</b>	<b>29</b>	<b>8</b>	<b>28%</b>	<b>18</b>	<b>62%</b>	<b>1</b>	<b>4</b>	<b>5</b>	<b>11</b>	<b>55</b>	<b>66</b>	<b>72</b>	<b>81</b>	<b>159</b>	<b>251</b>	<b>55</b>	<b>56.17</b>	<b>10</b>	<b>11</b>	<b>3</b>	<b>10%</b>
493	I	Asthma	159	72	45%	69	43%	1	5	10	16	26	34	90	160	365	365	26	86.83	5		18	11%
493.9	I	Asthma, unspecified	119	54	45%	53	45%	3	7	11	16	29	38	94	253	365	365	29	95.40	5		12	10%
<b>493.90</b>		<b>Asthma, unspecified</b>	<b>83</b>	<b>39</b>	<b>47%</b>	<b>38</b>	<b>46%</b>	<b>4</b>	<b>10</b>	<b>13</b>	<b>19</b>	<b>34</b>	<b>48</b>	<b>118</b>	<b>330</b>	<b>365</b>	<b>365</b>	<b>34</b>	<b>110.21</b>	<b>5</b>		<b>6</b>	<b>7%</b>
506	I	Respiratory conditions due to chemical fumes and vapors	351	204	58%	124	35%	1	1	2	3	4	5	6	14	27	355	4	18.45	4		23	7%
<b>506.0</b>		<b>Bronchitis and pneumonitis due to fumes and vapors</b>	<b>193</b>	<b>106</b>	<b>55%</b>	<b>76</b>	<b>39%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>7</b>	<b>13</b>	<b>26</b>	<b>232</b>	<b>4</b>	<b>14.46</b>	<b>4</b>		<b>11</b>	<b>6%</b>
<b>506.3</b>		<b>Other acute and subacute respiratory conditions due to fumes and vap</b>	<b>92</b>	<b>54</b>	<b>59%</b>	<b>32</b>	<b>35%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>4</b>	<b>6</b>	<b>16</b>	<b>21</b>	<b>48</b>	<b>355</b>	<b>4</b>	<b>24.91</b>	<b>5</b>		<b>6</b>	<b>7%</b>
511	I	Pleurisy	60	23	38%	28	47%	9	31	40	54	62	67	76	87	93	230	62	63.07	4		9	15%
<b>511.8</b>		<b>Other specified forms of effusion, except tuberculous</b>	<b>22</b>	<b>4</b>	<b>18%</b>	<b>15</b>	<b>68%</b>	<b>9</b>	<b>26</b>	<b>44</b>	<b>62</b>	<b>67</b>	<b>78</b>	<b>87</b>	<b>88</b>	<b>93</b>	<b>138</b>	<b>67</b>	<b>64.87</b>	<b>4</b>		<b>3</b>	<b>14%</b>
512	I	Pneumothorax	39		0%	36	92%	10	16	18	20	25	32	39	46	70	256	25	37.39	29		3	8%
<b>512.8</b>		<b>Other spontaneous pneumothorax</b>	<b>38</b>		<b>0%</b>	<b>35</b>	<b>92%</b>	<b>10</b>	<b>16</b>	<b>18</b>	<b>20</b>	<b>25</b>	<b>32</b>	<b>38</b>	<b>41</b>	<b>55</b>	<b>256</b>	<b>25</b>	<b>36.46</b>	<b>32</b>		<b>3</b>	<b>8%</b>
518	I	Other diseases of lung	58	8	14%	38	66%	15	42	50	62	78	106	112	145	271	365	78	108.45	16		12	21%
<b>518.0</b>		<b>Pulmonary collapse</b>	<b>34</b>	<b>2</b>	<b>6%</b>	<b>27</b>	<b>79%</b>	<b>15</b>	<b>38</b>	<b>54</b>	<b>62</b>	<b>80</b>	<b>108</b>	<b>111</b>	<b>145</b>	<b>312</b>	<b>365</b>	<b>80</b>	<b>110.33</b>	<b>16</b>	<b>21</b>	<b>5</b>	<b>15%</b>
524	I	Dentofacial anomalies, including malocclusion	67	26	39%	34	51%	1	2	4	5	6	12	14	77	134	259	6	38.00	2		7	10%
524.6	I	Temporomandibular joint disorders	46	16	35%	23	50%	1	2	3	5	6	7	53	84	156	259	6	43.48	6		7	15%
525	I	Other diseases and conditions of the teeth and supporting structures	167	117	70%	43	26%	1	1	2	2	2	3	4	5	12	35	2	5.28	2		7	4%
525.1	I	Loss of teeth due to trauma, extraction, or periodontal disease	89	56	63%	27	30%	1	1	2	2	3	4	4	7	29	35	3	6.78	2		6	7%
<b>525.11</b>		<b>Loss of teeth due to trauma</b>	<b>28</b>	<b>13</b>	<b>46%</b>	<b>15</b>	<b>54%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>12</b>	<b>29</b>	<b>30</b>	<b>3</b>	<b>7.67</b>	<b>5</b>			<b>0%</b>
550	I	Inguinal hernia	3,842	476	12%	3,157	82%	12	19	25	31	37	43	49	60	79	365	37	44.46	29	23	209	5%
550.1	I	Inguinal hernia, with obstruction, without mention of gangrene	78	2	3%	72	92%	11	18	24	31	40	45	48	59	67	365	40	44.65	34		4	5%
<b>550.10</b>		<b>Inguinal hernia, with obstruction, without mention of gangrene</b>	<b>66</b>	<b>2</b>	<b>3%</b>	<b>60</b>	<b>91%</b>	<b>10</b>	<b>15</b>	<b>20</b>	<b>29</b>	<b>35</b>	<b>41</b>	<b>46</b>	<b>55</b>	<b>65</b>	<b>365</b>	<b>35</b>	<b>42.65</b>	<b>34</b>		<b>4</b>	<b>6%</b>
550.9	I	Inguinal hernia, without mention of obstruction or gangrene	3,754	474	13%	3,077	82%	12	19	25	31	36	43	49	60	80	365	36	44.45	29		203	5%
<b>550.90</b>		<b>Inguinal hernia, without mention of obstruction or gangrene</b>	<b>3,438</b>	<b>445</b>	<b>13%</b>	<b>2,808</b>	<b>82%</b>	<b>11</b>	<b>19</b>	<b>25</b>	<b>31</b>	<b>35</b>	<b>43</b>	<b>49</b>	<b>60</b>	<b>78</b>	<b>365</b>	<b>35</b>	<b>43.56</b>	<b>29</b>		<b>185</b>	<b>5%</b>
<b>550.91</b>		<b>Inguinal hernia, without mention of obstruction or gangrene</b>	<b>139</b>	<b>13</b>	<b>9%</b>	<b>118</b>	<b>85%</b>	<b>15</b>	<b>22</b>	<b>30</b>	<b>36</b>	<b>41</b>	<b>46</b>	<b>53</b>	<b>75</b>	<b>92</b>	<b>365</b>	<b>41</b>	<b>53.07</b>	<b>29</b>		<b>8</b>	<b>6%</b>
<b>550.92</b>		<b>Inguinal hernia, without mention of obstruction or gangrene</b>	<b>174</b>	<b>16</b>	<b>9%</b>	<b>148</b>	<b>85%</b>	<b>16</b>	<b>20</b>	<b>28</b>	<b>34</b>	<b>42</b>	<b>48</b>	<b>57</b>	<b>71</b>	<b>99</b>	<b>365</b>	<b>42</b>	<b>54.34</b>	<b>29</b>		<b>10</b>	<b>6%</b>
552	I	Other hernia of abdominal cavity, with obstruction, but without mention of ga	126	8	6%	114	90%	14	22	27	32	38	44	49	58	74	224	38	45.54	11		4	3%
<b>552.1</b>		<b>Umbilical hernia with obstruction</b>	<b>76</b>	<b>6</b>	<b>8%</b>	<b>69</b>	<b>91%</b>	<b>13</b>	<b>19</b>	<b>24</b>	<b>30</b>	<b>35</b>	<b>42</b>	<b>48</b>	<b>58</b>	<b>74</b>	<b>198</b>	<b>35</b>	<b>42.04</b>	<b>24</b>		<b>1</b>	<b>1%</b>
552.2	I	Ventral hernia with obstruction	39	2	5%	35	90%	10	25	28	32	39	44	49	56	91	224	39	49.03	39		2	5%
<b>552.21</b>		<b>Incisional, with obstruction</b>	<b>18</b>	<b>1</b>	<b>6%</b>	<b>16</b>	<b>89%</b>	<b>25</b>	<b>31</b>	<b>38</b>	<b>41</b>	<b>44</b>	<b>49</b>	<b>56</b>	<b>58</b>	<b>141</b>	<b>166</b>	<b>44</b>	<b>54.69</b>	<b>45</b>		<b>1</b>	<b>6%</b>
553	I	Other hernia of abdominal cavity without mention of obstruction or gangrene	1,158	155	13%	935	81%	11	19	26	32	40	46	52	62	86	365	40	50.33	21		68	6%
553.0	I	Femoral hernia	30	4	13%	25	83%	9	18	25	30	39	42	47	52	112	176	39	45.44	38		1	3%
<b>553.00</b>		<b>Unilateral or unspecified (not specified as recurrent)</b>	<b>28</b>	<b>4</b>	<b>14%</b>	<b>23</b>	<b>82%</b>	<b>9</b>	<b>18</b>	<b>25</b>	<b>33</b>	<b>39</b>	<b>42</b>	<b>47</b>	<b>52</b>	<b>82</b>	<b>176</b>	<b>39</b>	<b>43.61</b>	<b>38</b>		<b>1</b>	<b>4%</b>
553.1	I	Umbilical hernia	627	85	14%	515	82%	11	18	24	31	36	43	48	56	81	365	36	46.46	32		27	4%
553.2	I	Ventral hernia	376	38	10%	309	82%	13	22	32	39	45	50	59	74	104	365	45	59.34	35	20	29	8%
<b>553.20</b>		<b>Ventral, unspecified</b>	<b>214</b>	<b>26</b>	<b>12%</b>	<b>170</b>	<b>79%</b>	<b>9</b>	<b>19</b>	<b>28</b>	<b>34</b>	<b>41</b>	<b>47</b>	<b>54</b>	<b>61</b>	<b>78</b>	<b>365</b>	<b>41</b>	<b>48.65</b>	<b>40</b>		<b>18</b>	<b>8%</b>
<b>553.21</b>		<b>Incisional</b>	<b>102</b>	<b>5</b>	<b>5%</b>	<b>89</b>	<b>87%</b>	<b>17</b>	<b>31</b>	<b>39</b>	<b>46</b>	<b>50</b>	<b>58</b>	<b>71</b>	<b>93</b>	<b>179</b>	<b>365</b>	<b>50</b>	<b>76.25</b>	<b>46</b>		<b>8</b>	<b>8%</b>
<b>553.29</b>		<b>Other</b>	<b>60</b>	<b>7</b>	<b>12%</b>	<b>50</b>	<b>83%</b>	<b>16</b>	<b>27</b>	<b>33</b>	<b>41</b>	<b>46</b>	<b>58</b>	<b>73</b>	<b>83</b>	<b>131</b>	<b>365</b>	<b>46</b>	<b>65.60</b>	<b>42</b>		<b>3</b>	<b>5%</b>
<b>553.8</b>		<b>Hernia of other specified sites</b>	<b>21</b>	<b>4</b>	<b>19%</b>	<b>15</b>	<b>71%</b>	<b>8</b>	<b>8</b>	<b>17</b>	<b>19</b>	<b>37</b>	<b>47</b>	<b>72</b>	<b>78</b>	<b>125</b>	<b>180</b>	<b>37</b>	<b>51.60</b>	<b>16</b>	<b>28</b>	<b>2</b>	<b>10%</b>
<b>553.9</b>		<b>Hernia of unspecified site</b>	<b>102</b>	<b>24</b>	<b>24%</b>	<b>69</b>	<b>68%</b>	<b>10</b>	<b>19</b>	<b>22</b>	<b>29</b>	<b>37</b>	<b>45</b>	<b>49</b>	<b>60</b>	<b>84</b>	<b>124</b>	<b>37</b>	<b>40.22</b>	<b>16</b>	<b>30</b>	<b>9</b>	<b>9%</b>
599	I	Other disorders of urethra and urinary tract	54	12	22%	34	63%	2	3	6	10	19	57	171	365	365	365	19	116.94	3		8	15%
<b>599.7</b>		<b>Hematuria</b>	<b>38</b>	<b>12</b>	<b>32%</b>	<b>24</b>	<b>63%</b>	<b>1</b>	<b>2</b>	<b>4</b>	<b>6</b>	<b>9</b>	<b>16</b>	<b>28</b>	<b>57</b>	<b>108</b>	<b>365</b>	<b>9</b>	<b>41.79</b>	<b>8</b>		<b>2</b>	<b>5%</b>
603	I	Hydrocele	47	10	21%	30	64%	2	5	9	23	36	41	48	73	124	271	36	51.63	21		7	15%
<b>603.9</b>		<b>Hydrocele, unspecified</b>	<b>44</b>	<b>8</b>	<b>18%</b>	<b>29</b>	<b>66%</b>	<b>2</b>	<b>5</b>	<b>9</b>	<b>23</b>	<b>36</b>	<b>41</b>	<b>56</b>	<b>84</b>	<b>170</b>	<b>271</b>	<b>36</b>	<b>52.07</b>	<b>21</b>		<b>7</b>	<b>16%</b>
604	I	Orchitis and epididymitis	129	60	47%	60	47%	1	3	5	5	6	7	11	18	28	76	6	12.30	10		9	7%
604.9	I	Other orchitis, epididymitis, and epididymo-orchitis, without mention of abscess	128	60	47%	59	46%	1	3	5	5	6	7	12	18	33	76	6	12.41	10		9	7%
<b>604.90</b>		<b>Orchitis and epididymitis, unspecified</b>	<b>108</b>	<b>51</b>	<b>47%</b>	<b>49</b>	<b>45%</b>	<b>1</b>	<b>3</b>	<b>5</b>	<b>6</b>	<b>6</b>	<b>7</b>	<b>12</b>	<b>20</b>	<b>37</b>	<b>76</b>	<b>6</b>	<b>13.22</b>	<b>6</b>		<b>8</b>	<b>7%</b>
608	I	Other disorders of male genital organs	66	22	33%	34	52%	1	2	5	9	11	19	23	29	39	316	11	27.41	8		10	15%
681	I	Cellulitis and abscess of finger and toe	1,610	1,118	69%	449	28%	1	1	2	3	4	6	9	14	26	365	4	11.91	9		43	3%
681.0	I	Finger	1,473	1,035	70%	398	27%																

### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
681.01		Felon	42	14	33%	24	57%	3	5	10	10	13	17	25	34	57	90	13	22.17	17		4	10%
681.02		Onychia and paronychia of finger	377	282	75%	85	23%	1	1	2	2	4	5	6	11	17	146	4	8.48	4		10	3%
681.1	I	Toe	103	55	53%	47	46%	1	2	3	4	5	7	12	19	36	82	5	12.13	10		1	1%
681.10		Cellulitis and abscess, unspecified	64	31	48%	33	52%	1	2	2	4	5	10	19	26	42	82	5	15.06	10			0%
682	I	Other cellulitis and abscess	3,112	1,655	53%	1,325	43%	1	2	3	4	6	11	16	24	44	365	6	19.06	15	6	132	4%
682.2		Trunk	48	23	48%	22	46%	2	3	6	8	15	22	35	88	118	195	15	39.45	18		3	6%
682.3		Upper arm and forearm	680	416	61%	238	35%	1	2	2	4	5	7	11	18	37	267	5	15.65	6		26	4%
682.4		Hand, except fingers and thumb	826	499	60%	293	35%	1	2	2	3	5	6	11	16	30	364	5	14.18	6		34	4%
682.6		Leg, except foot	1,071	489	46%	535	50%	2	3	4	6	10	14	20	29	50	365	10	21.24	9		47	4%
682.7		Foot, except toes	324	121	37%	187	58%	2	3	4	5	10	14	22	33	69	365	10	24.24	14		16	5%
682.9		Unspecified site	85	58	68%	23	27%	1	1	1	2	3	3	5	6	10	26	3	4.91	15		4	5%
686	I	Other local infections of skin and subcutaneous tissue	290	195	67%	83	29%	1	2	3	4	6	9	13	20	34	365	6	21.08	7		12	4%
686.9		Unspecified local infection of skin and subcutaneous tissue	239	158	66%	70	29%	1	1	2	3	5	9	10	14	24	279	5	14.54	7	7	11	5%
691	I	Atopic dermatitis and related conditions	64	37	58%	24	38%	1	2	2	2	2	5	6	12	44	343	2	22.04	3	2	3	5%
691.8		Other atopic dermatitis and related conditions	64	37	58%	24	38%	1	2	2	2	2	5	6	12	44	343	2	22.04	5		3	5%
692	I	Contact dermatitis and other eczema	6,940	5,158	74%	1,467	21%	1	1	2	2	3	4	5	6	15	365	3	11.60	3	3	315	5%
692.0		Due to detergents	110	88	80%	20	18%	1	1	1	2	2	2	3	4	5	8	2	2.65	2		2	2%
692.1		Due to oils and greases	114	84	74%	25	22%	1	1	2	3	3	3	6	12	19	69	3	9.72	3		5	4%
692.2		Due to solvents	162	103	64%	45	28%	1	2	2	3	3	4	6	8	15	98	3	9.00	4	2	14	9%
692.4		Due to other chemical products	1,228	929	76%	241	20%	1	1	2	3	3	5	8	16	38	365	3	23.64	3	3	58	5%
692.6		Due to plants [except food]	2,378	1,786	75%	489	21%	1	1	2	2	2	3	4	4	6	365	2	4.99	2		103	4%
692.8	I	Due to other specified agents	87	61	70%	21	24%	1	1	2	2	3	4	4	5	14	31	3	5.62	3	1	5	6%
692.89		Other	78	55	71%	18	23%	1	1	2	2	3	4	4	8	23	31	3	6.17	3		5	6%
692.9		Unspecified cause	2,773	2,044	74%	604	22%	1	1	2	2	3	4	5	8	19	365	3	11.93	3		125	5%
703	I	Diseases of nail	86	54	63%	29	34%	2	3	4	6	13	16	23	31	40	232	13	22.52	23	2	3	3%
706	I	Diseases of sebaceous glands	52	32	62%	19	37%	2	3	4	13	14	19	58	66	171	365	14	50.11	3	8	1	2%
706.2		Sebaceous cyst	50	32	64%	17	34%	2	3	6	13	14	19	21	65	108	171	14	34.35	8	8	1	2%
707	I	Chronic ulcer of skin	101	31	31%	60	59%	4	16	33	58	97	117	161	276	365	365	97	128.87	8		10	10%
707.1	I	Ulcer of lower limbs, except decubitus	84	23	27%	53	63%	4	13	30	58	97	115	157	186	321	365	97	120.64	16		8	10%
707.15		Ulcer of other part of foot	25	6	24%	19	76%	2	4	25	66	139	163	176	307	321	365	139	135.47	66			0%
708	I	Urticaria	170	128	75%	33	19%	1	1	1	1	2	2	3	4	6	365	2	13.30	4	1	9	5%
708.0		Allergic urticaria	84	61	73%	19	23%	1	1	1	1	2	2	2	4	6	8	2	2.47	4		4	5%
709	I	Other disorders of skin and subcutaneous tissue	224	116	52%	92	41%	2	4	6	10	14	18	30	75	162	365	14	49.96	10		16	7%
709.2		Scar conditions and fibrosis of skin	66	15	23%	44	67%	3	6	9	14	18	33	78	159	208	365	18	71.52	24		7	11%
709.4		Foreign body granuloma of skin and subcutaneous tissue	124	85	69%	33	27%	2	3	4	6	10	14	18	21	36	365	10	25.00	6		6	5%
711	I	Arthropathy associated with infections	30	3	10%	27	90%	7	9	15	18	23	29	32	74	365	365	23	68.89	28			0%
711.0	I	Pyogenic arthritis	26	2	8%	24	92%	7	9	14	19	23	29	32	74	365	365	23	68.21	19			0%
715	I	Osteoarthritis and allied disorders	2,073	245	12%	1,276	62%	14	39	63	93	124	167	230	333	365	365	124	161.05	27		552	27%
715.1	I	Osteoarthritis, localized, primary	576	67	12%	369	64%	18	44	69	95	125	165	228	306	365	365	125	160.43	14		140	24%
715.11		Osteoarthritis, localized, primary	331	23	7%	226	68%	33	54	78	100	137	185	249	321	365	365	137	170.78	14		82	25%
715.14		Osteoarthritis, localized, primary	32	9	28%	19	59%	2	9	33	63	89	102	222	365	365	365	89	143.05	14		4	13%
715.16		Osteoarthritis, localized, primary	117	22	19%	67	57%	13	28	70	88	107	125	159	259	365	365	107	138.24	14		28	24%
715.17		Osteoarthritis, localized, primary	27	1	4%	19	70%	10	35	60	84	98	137	187	228	338	365	98	137.26	14		7	26%
715.18		Osteoarthritis, localized, primary	32	3	9%	17	53%	9	39	55	150	232	295	365	365	365	365	232	202.88	14		12	38%
715.2	I	Osteoarthritis, localized, secondary	97	12	12%	50	52%	28	55	76	138	198	241	296	365	365	365	198	200.94	17		35	36%
715.26		Osteoarthritis, localized, secondary	28	4	14%	17	61%	41	62	123	136	186	252	287	365	365	365	186	198.76	17		7	25%
715.3	I	Osteoarthritis, localized, not specified whether primary or secondary	166	12	7%	102	61%	17	46	80	104	156	235	291	365	365	365	156	183.92	13		52	31%
715.31		Osteoarthritis, localized, not specified whether primary or secondary	81	4	5%	56	69%	18	48	84	107	156	238	318	365	365	365	156	188.45	13		21	26%
715.36		Osteoarthritis, localized, not specified whether primary or secondary	64	6	9%	38	59%	14	30	49	92	110	169	216	325	365	365	110	157.53	13		20	31%
715.9	I	Osteoarthritis, unspecified whether generalized or localized	1,217	149	12%	749	62%	13	35	59	89	116	160	214	328	365	365	116	156.46	27		319	26%
715.91		Osteoarthritis, unspecified whether generalized or localized	249	34	14%	158	63%	19	41	64	86	114	164	216	280	365	365	114	152.20	27		57	23%



**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
715.93		Osteoarthritis, unspecified whether generalized or localized	40	9	23%	24	60%	3	26	56	89	113	122	330	365	365	365	113	166.58	27		7	18%
715.94		Osteoarthritis, unspecified whether generalized or localized	75	22	29%	41	55%	18	27	68	96	113	170	276	365	365	365	113	165.37	27		12	16%
715.95		Osteoarthritis, unspecified whether generalized or localized	38	3	8%	20	53%	3	13	30	54	101	111	154	205	365	365	101	135.35	27		15	39%
715.96		Osteoarthritis, unspecified whether generalized or localized	738	74	10%	462	63%	9	35	57	90	120	161	214	348	365	365	120	158.05	27		202	27%
715.97		Osteoarthritis, unspecified whether generalized or localized	60	5	8%	33	55%	16	66	80	124	144	180	235	317	365	365	144	172.88	27		22	37%
716	I	Other and unspecified arthropathies	1,141	152	13%	647	57%	11	40	69	102	156	221	310	365	365	365	156	182.31	20		342	30%
716.1	I	Traumatic arthropathy	539	42	8%	323	60%	19	53	87	132	187	245	341	365	365	365	187	199.02	23		174	32%
716.11		Traumatic arthropathy	85	7	8%	60	71%	19	60	89	113	156	183	229	365	365	365	156	178.85	23		18	21%
716.13		Traumatic arthropathy	63	7	11%	36	57%	66	104	117	235	245	334	365	365	365	365	245	237.25	23		20	32%
716.14		Traumatic arthropathy	43	8	19%	31	72%	4	6	13	34	60	72	115	240	350	365	60	108.97	23		4	9%
716.16		Traumatic arthropathy	131	9	7%	71	54%	42	60	83	138	181	267	341	365	365	365	181	202.23	23		51	39%
716.17		Traumatic arthropathy	173	9	5%	101	58%	29	68	134	192	233	309	365	365	365	365	233	225.50	23		63	36%
716.8	I	Other specified arthropathy	50	10	20%	21	42%	6	26	58	95	125	156	286	365	365	365	125	165.14	10		19	38%
716.81		Other specified arthropathy	34	7	21%	16	47%	6	30	65	125	132	239	288	365	365	365	132	182.00	10		11	32%
716.9	I	Arthropathy, unspecified	543	98	18%	300	55%	7	25	51	88	121	184	275	365	365	365	121	165.91	25		145	27%
716.91		Arthropathy, unspecified	137	28	20%	83	61%	5	17	34	55	84	124	223	305	365	365	84	141.52	25		26	19%
716.93		Arthropathy, unspecified	34	8	24%	17	50%	3	14	89	99	120	123	181	228	257	365	120	131.00	25		9	26%
716.94		Arthropathy, unspecified	80	21	26%	38	48%	14	50	88	94	96	140	250	337	365	365	96	158.18	25		21	26%
716.96		Arthropathy, unspecified	143	19	13%	82	57%	11	32	51	59	97	155	229	361	365	365	97	153.94	25		42	29%
716.97		Arthropathy, unspecified	87	8	9%	55	63%	17	40	103	166	210	297	365	365	365	365	210	215.20	25		24	28%
717	I	Internal derangement of knee	1,256	282	22%	808	64%	5	18	34	49	70	92	129	195	339	365	70	110.91	45		166	13%
717.1		Derangement of anterior horn of medial meniscus	26	6	23%	18	69%	20	31	42	62	89	94	114	181	365	365	89	124.50	76		2	8%
717.2		Derangement of posterior horn of medial meniscus	69	8	12%	45	65%	28	36	55	84	105	134	202	262	365	365	105	144.24	45		16	23%
717.3		Other and unspecified derangement of medial meniscus	46	18	39%	24	52%	1	7	17	25	41	54	66	113	157	365	41	67.58	25		4	9%
717.5		Derangement of meniscus, not elsewhere classified	52	10	19%	29	56%	18	42	56	73	93	114	157	208	365	365	93	132.97	28		13	25%
717.6		Loose body in knee	115	26	23%	75	65%	11	20	35	45	56	77	96	121	264	365	56	91.16	25		14	12%
717.7		Chondromalacia of patella	766	165	22%	513	67%	4	17	33	49	71	93	139	198	342	365	71	112.50	30		88	11%
717.8	I	Other internal derangement of knee	159	44	28%	92	58%	4	16	25	47	52	83	112	213	307	365	52	105.08	28		23	14%
717.83		Old disruption of anterior cruciate ligament	49	6	12%	35	71%	14	18	47	61	92	105	150	235	365	365	92	130.17	31		8	16%
717.89		Other	75	32	43%	38	51%	3	7	16	21	41	52	77	193	217	365	41	82.13	18		5	7%
718	I	Other derangement of joint	1,271	207	16%	807	63%	10	28	53	78	111	159	235	322	365	365	111	155.04	8		257	20%
718.0	I	Articular cartilage disorder	109	10	9%	78	72%	19	44	56	74	125	188	264	311	365	365	125	166.14	10		21	19%
718.01		Articular cartilage disorder	63	5	8%	48	76%	19	53	59	83	117	188	256	304	365	365	117	160.56	10		10	16%
718.1	I	Loose body in joint	82	22	27%	45	55%	15	26	34	55	87	108	155	234	365	365	87	128.09	18		15	18%
718.12		Loose body in joint	39	12	31%	23	59%	15	21	26	32	44	56	108	155	234	365	44	93.00	18		4	10%
718.17		Loose body in joint	29	3	10%	15	52%	29	34	62	87	111	116	214	274	365	365	111	150.80	18		11	38%
718.3	I	Recurrent dislocation of joint	66	16	24%	41	62%	3	21	43	53	91	133	174	210	303	365	91	122.49	8		9	14%
718.31		Recurrent dislocation of joint	41	12	29%	24	59%	2	3	27	43	52	106	133	174	240	365	52	98.42	8		5	12%
718.4	I	Contracture of joint	142	24	17%	91	64%	10	19	52	88	112	175	261	365	365	365	112	163.14	16		27	19%
718.44		Contracture of joint	85	19	22%	57	67%	6	16	21	48	77	112	151	209	365	365	77	119.77	16		9	11%
718.5	I	Ankylosis of joint	277	27	10%	188	68%	8	31	53	73	96	128	194	275	365	365	96	142.85	19		62	22%
718.54		Ankylosis of joint	195	24	12%	149	76%	6	23	46	62	82	107	142	235	328	365	82	119.83	19		22	11%
718.56		Ankylosis of joint	46		0%	24	52%	104	118	206	263	292	365	365	365	365	365	292	267.63	19		22	48%
718.8	I	Other joint derangement, not elsewhere classified	585	107	18%	357	61%	9	28	54	89	129	179	264	365	365	365	129	164.51	14		121	21%
718.81		Other joint derangement, not elsewhere classified	214	21	10%	140	65%	35	64	96	123	163	227	301	365	365	365	163	195.28	14		53	25%
718.83		Other joint derangement, not elsewhere classified	111	19	17%	67	60%	8	24	47	84	178	223	303	365	365	365	178	178.46	14		25	23%
718.84		Other joint derangement, not elsewhere classified	49	30	61%	19	39%	1	1	2	4	5	16	30	45	132	365	5	39.37	14			0%
718.86		Other joint derangement, not elsewhere classified	43	15	35%	22	51%	15	20	20	28	41	60	73	272	365	365	41	106.05	14		6	14%
718.87		Other joint derangement, not elsewhere classified	153	15	10%	103	67%	13	31	58	98	126	151	224	318	365	365	126	157.67	14		35	23%
719	I	Other and unspecified disorders of joint	757	254	34%	427	56%	2	5	10	16	30	46	74	128	258	365	30	74.95	11		76	10%
719.0	I	Effusion of joint	434	182	42%	224	52%	1	2	5	8	14	22	37	59	115	365	14	43.04	11		28	6%

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
719.02		<b>Effusion of joint</b>	<b>40</b>	<b>22</b>	<b>55%</b>	<b>15</b>	<b>38%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>6</b>	<b>14</b>	<b>20</b>	<b>24</b>	<b>59</b>	<b>117</b>	<b>6</b>	<b>20.40</b>	<b>11</b>		<b>3</b>	<b>8%</b>
719.06		<b>Effusion of joint</b>	<b>319</b>	<b>126</b>	<b>39%</b>	<b>175</b>	<b>55%</b>	<b>1</b>	<b>2</b>	<b>4</b>	<b>8</b>	<b>12</b>	<b>19</b>	<b>29</b>	<b>46</b>	<b>83</b>	<b>365</b>	<b>12</b>	<b>34.58</b>	<b>11</b>		<b>18</b>	<b>6%</b>
719.1	I	Hemarthrosis	42	7	17%	31	74%	6	11	12	18	34	74	95	151	210	365	34	82.68	13		4	10%
719.16		<b>Hemarthrosis</b>	<b>27</b>	<b>2</b>	<b>7%</b>	<b>23</b>	<b>85%</b>	<b>11</b>	<b>12</b>	<b>13</b>	<b>34</b>	<b>74</b>	<b>95</b>	<b>151</b>	<b>167</b>	<b>241</b>	<b>365</b>	<b>74</b>	<b>104.48</b>	<b>13</b>		<b>2</b>	<b>7%</b>
719.2	I	Villonodular synovitis	105	16	15%	70	67%	14	32	43	59	77	102	163	286	365	365	77	135.07	33		19	18%
719.21		<b>Villonodular synovitis</b>	<b>18</b>	<b>1</b>	<b>6%</b>	<b>15</b>	<b>83%</b>	<b>33</b>	<b>54</b>	<b>66</b>	<b>72</b>	<b>91</b>	<b>92</b>	<b>206</b>	<b>321</b>	<b>365</b>	<b>365</b>	<b>91</b>	<b>159.53</b>	<b>33</b>		<b>2</b>	<b>11%</b>
719.26		<b>Villonodular synovitis</b>	<b>52</b>	<b>7</b>	<b>13%</b>	<b>35</b>	<b>67%</b>	<b>14</b>	<b>19</b>	<b>41</b>	<b>46</b>	<b>71</b>	<b>94</b>	<b>163</b>	<b>278</b>	<b>365</b>	<b>365</b>	<b>71</b>	<b>131.57</b>	<b>33</b>		<b>10</b>	<b>19%</b>
719.4	I	Pain in joint	107	33	31%	63	59%	2	3	8	14	37	47	89	124	203	365	37	72.05	11		11	10%
719.46		<b>Pain in joint</b>	<b>55</b>	<b>13</b>	<b>24%</b>	<b>34</b>	<b>62%</b>	<b>2</b>	<b>2</b>	<b>25</b>	<b>39</b>	<b>42</b>	<b>77</b>	<b>130</b>	<b>183</b>	<b>325</b>	<b>365</b>	<b>42</b>	<b>98.97</b>	<b>11</b>		<b>8</b>	<b>15%</b>
719.8	I	Other specified disorders of joint	44	8	18%	25	57%	31	42	47	69	131	156	213	258	365	365	131	153.68	15		11	25%
720	I	Ankylosing spondylitis and other inflammatory spondylopathies	144	38	26%	91	63%	2	3	4	5	13	30	51	101	264	365	13	67.35	15		15	10%
720.2		<b>Sacroiliitis, not elsewhere classified</b>	<b>139</b>	<b>37</b>	<b>27%</b>	<b>87</b>	<b>63%</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>12</b>	<b>30</b>	<b>51</b>	<b>101</b>	<b>264</b>	<b>365</b>	<b>12</b>	<b>65.60</b>	<b>12</b>		<b>15</b>	<b>11%</b>
721	I	Spondylosis and allied disorders	990	164	17%	450	45%	4	17	41	80	120	172	254	365	365	365	120	162.00	12		376	38%
721.0		<b>Cervical spondylosis without myelopathy</b>	<b>418</b>	<b>68</b>	<b>16%</b>	<b>193</b>	<b>46%</b>	<b>4</b>	<b>20</b>	<b>46</b>	<b>73</b>	<b>123</b>	<b>172</b>	<b>277</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>123</b>	<b>162.88</b>	<b>12</b>		<b>157</b>	<b>38%</b>
721.1		<b>Cervical spondylosis with myelopathy</b>	<b>46</b>	<b>3</b>	<b>7%</b>	<b>21</b>	<b>46%</b>	<b>56</b>	<b>86</b>	<b>93</b>	<b>108</b>	<b>157</b>	<b>211</b>	<b>249</b>	<b>274</b>	<b>365</b>	<b>365</b>	<b>157</b>	<b>181.33</b>	<b>25</b>		<b>22</b>	<b>48%</b>
721.3		<b>Lumbosacral spondylosis without myelopathy</b>	<b>494</b>	<b>84</b>	<b>17%</b>	<b>221</b>	<b>45%</b>	<b>3</b>	<b>11</b>	<b>28</b>	<b>71</b>	<b>120</b>	<b>166</b>	<b>291</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>120</b>	<b>161.81</b>	<b>11</b>		<b>189</b>	<b>38%</b>
722	I	Intervertebral disc disorders	12,077	1,715	14%	6,806	56%	6	19	44	74	110	165	253	365	365	365	110	158.32	30	61	3,556	29%
722.0		<b>Displacement of cervical intervertebral disc without myelopathy</b>	<b>1,816</b>	<b>294</b>	<b>16%</b>	<b>1,007</b>	<b>55%</b>	<b>8</b>	<b>35</b>	<b>61</b>	<b>93</b>	<b>126</b>	<b>181</b>	<b>252</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>126</b>	<b>165.89</b>	<b>36</b>		<b>515</b>	<b>28%</b>
722.1	I	Displacement of thoracic or lumbar intervertebral disc without myelopathy	5,881	816	14%	3,587	61%	6	20	42	72	105	156	234	355	365	365	105	153.18	42		1,478	25%
722.10		<b>Lumbar intervertebral disc without myelopathy</b>	<b>5,714</b>	<b>787</b>	<b>14%</b>	<b>3,496</b>	<b>61%</b>	<b>6</b>	<b>21</b>	<b>43</b>	<b>72</b>	<b>105</b>	<b>156</b>	<b>233</b>	<b>355</b>	<b>365</b>	<b>365</b>	<b>105</b>	<b>153.31</b>	<b>29</b>		<b>1,431</b>	<b>25%</b>
722.11		<b>Thoracic intervertebral disc without myelopathy</b>	<b>167</b>	<b>29</b>	<b>17%</b>	<b>91</b>	<b>54%</b>	<b>2</b>	<b>7</b>	<b>13</b>	<b>28</b>	<b>86</b>	<b>141</b>	<b>285</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>86</b>	<b>148.34</b>	<b>19</b>		<b>47</b>	<b>28%</b>
722.2		<b>Displacement of intervertebral disc, site unspecified, without myelopathy</b>	<b>2,120</b>	<b>288</b>	<b>14%</b>	<b>1,115</b>	<b>53%</b>	<b>5</b>	<b>15</b>	<b>37</b>	<b>69</b>	<b>110</b>	<b>178</b>	<b>283</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>110</b>	<b>161.72</b>	<b>30</b>		<b>717</b>	<b>34%</b>
722.4		<b>Degeneration of cervical intervertebral disc</b>	<b>288</b>	<b>59</b>	<b>20%</b>	<b>138</b>	<b>48%</b>	<b>3</b>	<b>13</b>	<b>31</b>	<b>56</b>	<b>97</b>	<b>166</b>	<b>216</b>	<b>270</b>	<b>365</b>	<b>365</b>	<b>97</b>	<b>140.64</b>	<b>27</b>		<b>91</b>	<b>32%</b>
722.5	I	Degeneration of thoracic or lumbar intervertebral disc	1,671	239	14%	827	49%	5	14	33	64	109	173	310	365	365	365	109	162.24	21		605	36%
722.51		<b>Thoracic or thoracolumbar intervertebral disc</b>	<b>62</b>	<b>9</b>	<b>15%</b>	<b>36</b>	<b>58%</b>	<b>2</b>	<b>18</b>	<b>32</b>	<b>62</b>	<b>122</b>	<b>132</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>122</b>	<b>168.56</b>	<b>18</b>		<b>17</b>	<b>27%</b>
722.52		<b>Lumbar or lumbosacral intervertebral disc</b>	<b>1,609</b>	<b>230</b>	<b>14%</b>	<b>791</b>	<b>49%</b>	<b>5</b>	<b>14</b>	<b>33</b>	<b>66</b>	<b>108</b>	<b>173</b>	<b>309</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>108</b>	<b>161.95</b>	<b>17</b>		<b>588</b>	<b>37%</b>
722.7	I	Intervertebral disc disorder with myelopathy	80	12	15%	52	65%	2	4	20	46	96	134	206	331	365	365	96	135.50	15		16	20%
722.71		<b>Cervical region</b>	<b>22</b>	<b>2</b>	<b>9%</b>	<b>15</b>	<b>68%</b>	<b>1</b>	<b>2</b>	<b>46</b>	<b>62</b>	<b>106</b>	<b>166</b>	<b>331</b>	<b>332</b>	<b>365</b>	<b>365</b>	<b>106</b>	<b>164.13</b>	<b>15</b>		<b>5</b>	<b>23%</b>
722.73		<b>Lumbar region</b>	<b>53</b>	<b>10</b>	<b>19%</b>	<b>34</b>	<b>64%</b>	<b>2</b>	<b>3</b>	<b>11</b>	<b>37</b>	<b>49</b>	<b>122</b>	<b>164</b>	<b>227</b>	<b>365</b>	<b>365</b>	<b>49</b>	<b>121.59</b>	<b>23</b>		<b>9</b>	<b>17%</b>
722.8	I	Postlaminectomy syndrome	140		0%	41	29%	105	359	365	365	365	365	365	365	365	365	365	316.73	30		99	71%
722.83		<b>Lumbar region</b>	<b>140</b>		<b>0%</b>	<b>41</b>	<b>29%</b>	<b>105</b>	<b>359</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>316.73</b>	<b>30</b>		<b>99</b>	<b>71%</b>
722.9	I	Other and unspecified disc disorder	76	7	9%	36	47%	12	31	59	124	135	304	365	365	365	365	135	192.08	22		33	43%
722.93		<b>Lumbar region</b>	<b>59</b>	<b>2</b>	<b>3%</b>	<b>29</b>	<b>49%</b>	<b>14</b>	<b>33</b>	<b>98</b>	<b>135</b>	<b>185</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>185</b>	<b>214.14</b>	<b>22</b>		<b>28</b>	<b>47%</b>
723	I	Other disorders of cervical region	775	355	46%	338	44%	2	3	5	13	27	47	72	150	324	365	27	80.94	17		82	11%
723.0		<b>Spinal stenosis of cervical region</b>	<b>84</b>	<b>13</b>	<b>15%</b>	<b>40</b>	<b>48%</b>	<b>2</b>	<b>4</b>	<b>42</b>	<b>66</b>	<b>150</b>	<b>176</b>	<b>282</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>150</b>	<b>166.53</b>	<b>17</b>		<b>31</b>	<b>37%</b>
723.1		<b>Cervicalgia</b>	<b>79</b>	<b>41</b>	<b>52%</b>	<b>34</b>	<b>43%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>14</b>	<b>33</b>	<b>74</b>	<b>365</b>	<b>3</b>	<b>29.50</b>	<b>12</b>		<b>4</b>	<b>5%</b>
723.3		<b>Cervicobrachial syndrome (diffuse)</b>	<b>75</b>	<b>40</b>	<b>53%</b>	<b>32</b>	<b>43%</b>	<b>4</b>	<b>4</b>	<b>5</b>	<b>9</b>	<b>14</b>	<b>21</b>	<b>28</b>	<b>71</b>	<b>109</b>	<b>365</b>	<b>14</b>	<b>44.25</b>	<b>17</b>		<b>3</b>	<b>4%</b>
723.4		<b>Brachia neuritis or radiculitis NOS</b>	<b>495</b>	<b>243</b>	<b>49%</b>	<b>211</b>	<b>43%</b>	<b>2</b>	<b>4</b>	<b>6</b>	<b>17</b>	<b>33</b>	<b>53</b>	<b>75</b>	<b>135</b>	<b>276</b>	<b>365</b>	<b>33</b>	<b>80.21</b>	<b>26</b>		<b>41</b>	<b>8%</b>
724	I	Other and unspecified disorders of back	1,974	514	26%	1,042	53%	2	3	6	13	22	46	88	163	342	365	22	85.59	14		418	21%
724.0	I	Spinal stenosis, other than cervical	524	56	11%	234	45%	5	19	51	91	119	200	276	365	365	365	119	169.47	27		234	45%
724.02		<b>Lumbar region</b>	<b>523</b>	<b>56</b>	<b>11%</b>	<b>234</b>	<b>45%</b>	<b>5</b>	<b>19</b>	<b>51</b>	<b>91</b>	<b>119</b>	<b>200</b>	<b>276</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>119</b>	<b>169.47</b>	<b>39</b>		<b>233</b>	<b>45%</b>
724.2		<b>Lumbago</b>	<b>92</b>	<b>35</b>	<b>38%</b>	<b>48</b>	<b>52%</b>	<b>1</b>	<b>3</b>	<b>4</b>	<b>7</b>	<b>12</b>	<b>15</b>	<b>22</b>	<b>31</b>	<b>87</b>	<b>365</b>	<b>12</b>	<b>32.67</b>	<b>6</b>	<b>7</b>	<b>9</b>	<b>10%</b>
724.3		<b>Sciatica</b>	<b>461</b>	<b>155</b>	<b>34%</b>	<b>272</b>	<b>59%</b>	<b>2</b>	<b>3</b>	<b>6</b>	<b>11</b>	<b>17</b>	<b>31</b>	<b>45</b>	<b>77</b>	<b>159</b>	<b>365</b>	<b>17</b>	<b>54.72</b>	<b>8</b>	<b>17</b>	<b>34</b>	<b>7%</b>
724.4		<b>Thoracic or lumbosacral neuritis or radiculitis, unspecified</b>	<b>703</b>	<b>216</b>	<b>31%</b>	<b>384</b>	<b>55%</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>9</b>	<b>15</b>	<b>29</b>	<b>60</b>	<b>119</b>	<b>282</b>	<b>365</b>	<b>15</b>	<b>71.04</b>	<b>18</b>		<b>103</b>	<b>15%</b>
724.6		<b>Disorders of sacrum</b>	<b>66</b>	<b>15</b>	<b>23%</b>	<b>31</b>	<b>47%</b>	<b>3</b>	<b>5</b>	<b>10</b>	<b>12</b>	<b>32</b>	<b>73</b>	<b>101</b>	<b>126</b>	<b>337</b>	<b>365</b>	<b>32</b>	<b>88.65</b>	<b>15</b>		<b>20</b>	<b>30%</b>
724.7	I	Disorders of coccyx	30	9	30%																		

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
726.10		Disorders of bursae and tendons in shoulder region, unspecified	2,733	1,283	47%	1,225	45%	2	4	7	17	36	59	92	141	273	365	36	81.93	14		225	8%
726.11		Calcifying tendinitis of shoulder	124	68	55%	54	44%	1	2	3	5	7	17	32	74	156	365	7	45.67	7		2	2%
726.12		Bicipital tenosynovitis	488	295	60%	174	36%	1	2	3	4	10	19	28	54	126	365	10	42.33	10		19	4%
726.19		Other specified disorders	198	63	32%	105	53%	3	6	20	41	62	86	104	179	365	365	62	102.78	15		30	15%
726.2		Other affections of shoulder region, not elsewhere classified	2,519	423	17%	1,659	66%	13	33	55	81	110	149	208	305	365	365	110	151.17	46		437	17%
726.3	I	Enthesopathy of elbow region	5,725	3,453	60%	1,953	34%	1	3	4	8	18	31	53	85	178	365	18	57.55	16		319	6%
726.30		Enthesopathy of elbow, unspecified	72	41	57%	26	36%	1	4	8	18	33	72	98	127	207	365	33	77.31	8		5	7%
726.31		Medial epicondylitis	787	472	60%	261	33%	2	3	6	12	20	32	55	102	198	365	20	65.64	10		54	7%
726.32		Lateral epicondylitis	3,918	2,336	60%	1,359	35%	2	3	5	12	21	36	62	97	200	365	21	63.57	13	14	223	6%
726.33		Olecranon bursitis	929	593	64%	301	32%	1	2	2	4	5	9	18	30	54	300	5	21.14	5		35	4%
726.4		Enthesopathy of wrist and carpus	110	62	56%	42	38%	1	3	4	5	7	13	15	44	98	231	7	32.12	10		6	5%
726.5		Enthesopathy of hip region	232	103	44%	112	48%	1	2	3	5	9	18	47	86	263	365	9	64.33	6		17	7%
726.6	I	Enthesopathy of knee	1,062	570	54%	454	43%	1	2	3	4	6	11	20	30	53	365	6	25.82	13		38	4%
726.60		Enthesopathy of knee, unspecified	190	111	58%	73	38%	2	2	3	4	4	6	10	16	38	131	4	11.79	5		6	3%
726.64		Patellar tendinitis	182	91	50%	80	44%	1	3	4	5	8	13	22	34	70	365	8	31.33	6		11	6%
726.65		Prepatellar bursitis	611	321	53%	273	45%	1	2	3	4	6	12	22	32	54	365	6	27.05	9		17	3%
726.69		Other	41	24	59%	15	37%	2	2	3	4	6	9	15	20	79	352	6	35.73	5		2	5%
726.7	I	Enthesopathy of ankle and tarsus	577	235	41%	278	48%	2	3	7	14	28	51	84	142	314	365	28	82.08	20		64	11%
726.71		Achilles bursitis or tendinitis	229	101	44%	112	49%	2	3	5	7	11	18	33	52	127	365	11	45.52	8		16	7%
726.72		Tibialis tendinitis	156	50	32%	82	53%	3	11	25	42	68	88	132	211	335	365	68	112.40	24		24	15%
726.79		Other	140	52	37%	70	50%	2	4	12	29	51	74	146	283	365	365	51	112.97	11		18	13%
726.8		Other peripheral enthesopathies	38	20	53%	15	39%	1	2	5	5	7	19	177	235	365	365	7	102.67	6		3	8%
726.9	I	Unspecified enthesopathy	323	177	55%	110	34%	1	2	4	6	16	31	83	183	365	365	16	82.36	12		36	11%
726.90		Enthesopathy of unspecified site	245	159	65%	67	27%	1	1	2	3	5	6	11	21	77	365	5	32.96	5		19	8%
726.91		Exostosis of unspecified site	78	18	23%	43	55%	12	23	55	86	126	183	227	365	365	365	126	159.35	21		17	22%
727	I	Other disorders of synovium, tendon, and bursa	9,799	5,562	57%	3,684	38%	1	3	4	9	17	31	49	85	161	365	17	54.48	13		553	6%
727.0	I	Synovitis and tenosynovitis	8,043	4,894	61%	2,735	34%	1	2	3	6	12	20	33	56	118	365	12	42.61	9	19	414	5%
727.00		Synovitis and tenosynovitis, unspecified	293	123	42%	137	47%	2	10	22	34	47	61	97	139	271	365	47	89.49	9	20	33	11%
727.03		Trigger finger (acquired)	631	296	47%	301	48%	4	12	16	21	27	40	58	85	147	365	27	60.92	12		34	5%
727.04		Radial styloid tenosynovitis	973	517	53%	389	40%	2	3	8	19	30	44	65	110	200	365	30	68.14	7		67	7%
727.05		Other tenosynovitis or hand and wrist	5,026	3,313	66%	1,497	30%	1	2	3	4	6	10	17	35	72	365	6	28.57	6		216	4%
727.06		Tenosynovitis of foot and ankle	271	130	48%	121	45%	2	3	4	8	18	30	48	76	152	365	18	46.48	9		20	7%
727.09		Other	837	510	61%	283	34%	1	2	3	4	7	13	24	45	97	365	7	37.33	6		44	5%
727.4	I	Ganglion and cyst of synovium, tendon, and bursa	753	425	56%	271	36%	2	4	10	18	27	37	59	97	172	365	27	59.76	17	11	57	8%
727.40		Synovial cyst, unspecified	37	11	30%	21	57%	12	19	26	84	91	160	203	245	300	365	91	134.19	17		5	14%
727.41		Ganglion of joint	142	78	55%	56	39%	2	5	8	19	26	37	57	74	159	363	26	51.32	19		8	6%
727.42		Ganglion of tendon sheath	67	42	63%	21	31%	3	4	11	26	36	40	54	108	119	213	36	52.24	13		4	6%
727.43		Ganglion, unspecified	502	292	58%	170	34%	2	3	7	14	26	33	49	93	163	365	26	55.02	14		40	8%
727.6	I	Rupture of tendon, nontraumatic	732	175	24%	500	68%	4	13	30	50	69	94	114	157	268	365	69	99.62	34		57	8%
727.61		Complete rupture of rotator cuff	38	12	32%	22	58%	11	41	64	71	98	133	203	265	365	365	98	142.41	49	26	4	11%
727.62		Tendons of biceps (long head)	378	95	25%	265	70%	3	9	24	40	60	88	110	139	268	365	60	94.04	20		18	5%
727.63		Extensor tendons of hand and wrist	59	25	42%	29	49%	2	4	14	27	39	54	77	85	171	292	39	58.93	27		5	8%
727.64		Flexor tendons of hand and wrist	38	10	26%	27	71%	3	12	17	18	22	51	61	86	137	353	22	61.63	22		1	3%
727.65		Quadriceps tendon	41	3	7%	30	73%	13	50	80	95	129	162	229	256	365	365	129	166.17	62		8	20%
727.67		Achilles tendon	97	19	20%	71	73%	5	20	40	58	89	103	124	148	183	365	89	93.08	74		7	7%
727.68		Other tendons of foot and ankle	47	7	15%	30	64%	29	47	87	95	105	119	140	315	365	365	105	151.43	87		10	21%
727.8	I	Other disorders of synovium, tendon, and bursa	199	33	17%	151	76%	8	23	46	63	80	99	146	194	301	365	80	116.09	27		15	8%
727.83		Plica syndrome	125	14	11%	99	79%	12	31	52	69	91	134	167	235	365	365	91	131.22	27		12	10%
727.89		Other	41	9	22%	31	76%	3	25	44	47	63	75	88	110	170	365	63	87.81	32		1	2%
728	I	Disorders of muscle, ligament, and fascia	583	239	41%	303	52%	2	3	5	13	25	42	66	111	270	365	25	73.22	21	91	41	7%
728.6		Contracture of palmar fascia	24	2	8%	19	79%	2	20	45	66	90	118	135	193	365	365	90	122.58	20		3	13%

### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
728.7	I	Other fibromatoses	319	167	52%	135	42%	1	2	3	6	19	34	57	100	198	365	19	61.98	10		17	5%
<b>728.71</b>		<b>Plantar fascial fibromatosis</b>	<b>315</b>	<b>167</b>	<b>53%</b>	<b>131</b>	<b>42%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>6</b>	<b>18</b>	<b>34</b>	<b>53</b>	<b>94</b>	<b>189</b>	<b>365</b>	<b>18</b>	<b>57.46</b>	<b>14</b>		<b>17</b>	<b>5%</b>
728.8	I	Other disorders of muscle, ligament, and fascia	188	63	34%	114	61%	2	3	5	8	15	24	33	56	180	365	15	52.69	8		11	6%
<b>728.85</b>		<b>Spasm of muscle</b>	<b>126</b>	<b>47</b>	<b>37%</b>	<b>73</b>	<b>58%</b>	<b>1</b>	<b>2</b>	<b>4</b>	<b>5</b>	<b>8</b>	<b>15</b>	<b>24</b>	<b>31</b>	<b>66</b>	<b>365</b>	<b>8</b>	<b>34.21</b>	<b>6</b>		<b>6</b>	<b>5%</b>
729	I	Other disorders of soft tissues	718	347	48%	304	42%	2	3	5	8	14	26	46	93	239	365	14	62.61	7	25	67	9%
<b>729.1</b>		<b>Mylagia and myositis, unspecified</b>	<b>269</b>	<b>100</b>	<b>37%</b>	<b>135</b>	<b>50%</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>13</b>	<b>19</b>	<b>34</b>	<b>65</b>	<b>120</b>	<b>307</b>	<b>365</b>	<b>19</b>	<b>78.43</b>	<b>13</b>		<b>34</b>	<b>13%</b>
<b>729.2</b>		<b>Neuralgia, neuritis, and radiculitis, unspecified</b>	<b>170</b>	<b>67</b>	<b>39%</b>	<b>83</b>	<b>49%</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>9</b>	<b>15</b>	<b>27</b>	<b>79</b>	<b>133</b>	<b>259</b>	<b>365</b>	<b>15</b>	<b>71.80</b>	<b>13</b>		<b>20</b>	<b>12%</b>
<b>729.6</b>		<b>Residual foreign body in soft tissue</b>	<b>225</b>	<b>155</b>	<b>69%</b>	<b>62</b>	<b>28%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>14</b>	<b>19</b>	<b>30</b>	<b>88</b>	<b>4</b>	<b>11.19</b>	<b>4</b>		<b>8</b>	<b>4%</b>
730	I	Osteomyelitis, periostitis, and other infections involving bone	112	16	14%	78	70%	16	35	52	68	87	108	145	227	365	365	87	128.91	14		18	16%
730.0	I	Acute osteomyelitis	48	3	6%	39	81%	17	26	48	59	73	93	118	227	365	365	73	121.97	44		6	13%
<b>730.04</b>		<b>Acute osteomyelitis</b>	<b>18</b>	<b>1</b>	<b>6%</b>	<b>16</b>	<b>89%</b>	<b>13</b>	<b>19</b>	<b>20</b>	<b>44</b>	<b>53</b>	<b>73</b>	<b>98</b>	<b>161</b>	<b>330</b>	<b>365</b>	<b>53</b>	<b>98.63</b>	<b>44</b>		<b>1</b>	<b>6%</b>
730.2	I	Unspecified osteomyelitis	38	5	13%	26	68%	6	28	44	67	87	101	139	194	306	365	87	114.65	49		7	18%
732	I	Osteochondropathies	163	22	13%	106	65%	13	35	60	77	100	145	194	300	365	365	100	147.71	16		35	21%
<b>732.7</b>		<b>Osteochondritis dissecans</b>	<b>99</b>	<b>14</b>	<b>14%</b>	<b>67</b>	<b>68%</b>	<b>10</b>	<b>32</b>	<b>60</b>	<b>67</b>	<b>87</b>	<b>129</b>	<b>188</b>	<b>279</b>	<b>365</b>	<b>365</b>	<b>87</b>	<b>137.64</b>	<b>32</b>		<b>18</b>	<b>18%</b>
<b>732.9</b>		<b>Unspecified osteochondropathy</b>	<b>27</b>	<b>2</b>	<b>7%</b>	<b>19</b>	<b>70%</b>	<b>33</b>	<b>43</b>	<b>79</b>	<b>90</b>	<b>101</b>	<b>108</b>	<b>167</b>	<b>362</b>	<b>365</b>	<b>365</b>	<b>101</b>	<b>150.84</b>	<b>16</b>		<b>6</b>	<b>22%</b>
733	I	Other disorders of bone and cartilage	1,488	239	16%	941	63%	6	21	41	61	91	136	190	286	365	365	91	138.72	16		308	21%
733.1	I	Pathologic fracture	69	19	28%	38	55%	2	16	25	35	42	78	135	161	365	365	42	104.05	16		12	17%
733.2	I	Cyst of bone	23	3	13%	15	65%	2	8	34	38	81	99	194	198	365	365	81	126.93	14		5	22%
733.4	I	Aseptic necrosis of bone	70	1	1%	48	69%	29	65	82	105	124	148	246	365	365	365	124	171.71	92		21	30%
<b>733.42</b>		<b>Head and neck of femur</b>	<b>36</b>		<b>0%</b>	<b>27</b>	<b>75%</b>	<b>38</b>	<b>66</b>	<b>90</b>	<b>105</b>	<b>124</b>	<b>209</b>	<b>335</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>124</b>	<b>185.89</b>	<b>105</b>		<b>9</b>	<b>25%</b>
<b>733.6</b>		<b>Tietze's disease</b>	<b>230</b>	<b>127</b>	<b>55%</b>	<b>92</b>	<b>40%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>7</b>	<b>13</b>	<b>23</b>	<b>39</b>	<b>365</b>	<b>4</b>	<b>21.88</b>	<b>10</b>		<b>11</b>	<b>5%</b>
733.8	I	Malunion and nonunion of fracture	450	23	5%	295	66%	17	48	79	111	164	205	286	365	365	365	164	182.39	78		132	29%
<b>733.81</b>		<b>Malunion of fracture</b>	<b>80</b>	<b>3</b>	<b>4%</b>	<b>53</b>	<b>66%</b>	<b>22</b>	<b>49</b>	<b>79</b>	<b>101</b>	<b>160</b>	<b>224</b>	<b>310</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>160</b>	<b>185.75</b>	<b>86</b>		<b>24</b>	<b>30%</b>
<b>733.82</b>		<b>Nonunion of fracture</b>	<b>370</b>	<b>20</b>	<b>5%</b>	<b>242</b>	<b>65%</b>	<b>14</b>	<b>48</b>	<b>79</b>	<b>118</b>	<b>165</b>	<b>203</b>	<b>285</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>165</b>	<b>181.65</b>	<b>91</b>		<b>108</b>	<b>29%</b>
733.9	I	Other and unspecified disorders of bone and cartilage	642	65	10%	451	70%	16	33	48	60	86	119	174	244	365	365	86	133.33	16		126	20%
<b>733.90</b>		<b>Disorder of bone and cartilage, unspecified</b>	<b>38</b>	<b>3</b>	<b>8%</b>	<b>30</b>	<b>79%</b>	<b>20</b>	<b>33</b>	<b>42</b>	<b>53</b>	<b>75</b>	<b>109</b>	<b>181</b>	<b>251</b>	<b>365</b>	<b>365</b>	<b>75</b>	<b>135.43</b>	<b>42</b>		<b>5</b>	<b>13%</b>
<b>733.92</b>		<b>Chondromalacia</b>	<b>515</b>	<b>36</b>	<b>7%</b>	<b>369</b>	<b>72%</b>	<b>17</b>	<b>35</b>	<b>52</b>	<b>67</b>	<b>92</b>	<b>134</b>	<b>183</b>	<b>250</b>	<b>365</b>	<b>365</b>	<b>92</b>	<b>138.66</b>	<b>31</b>		<b>110</b>	<b>21%</b>
<b>733.99</b>		<b>Other</b>	<b>52</b>	<b>15</b>	<b>29%</b>	<b>32</b>	<b>62%</b>	<b>3</b>	<b>10</b>	<b>14</b>	<b>19</b>	<b>25</b>	<b>55</b>	<b>76</b>	<b>173</b>	<b>242</b>	<b>365</b>	<b>25</b>	<b>84.16</b>	<b>13</b>		<b>5</b>	<b>10%</b>
735	I	Acquired deformities of toe	27	10	37%	15	56%	3	4	7	27	33	36	46	72	365	365	33	73.73	12		2	7%
736	I	Other acquired deformities of limbs	545	297	54%	190	35%	2	4	9	27	47	74	125	179	365	365	47	99.67	16		58	11%
<b>736.1</b>		<b>Mallet finger</b>	<b>302</b>	<b>235</b>	<b>78%</b>	<b>57</b>	<b>19%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>8</b>	<b>38</b>	<b>69</b>	<b>143</b>	<b>4</b>	<b>21.40</b>	<b>5</b>		<b>10</b>	<b>3%</b>
736.2	I	Other acquired deformities of finger	118	53	45%	60	51%	2	5	13	27	45	49	86	157	244	365	45	86.42	28		5	4%
<b>736.21</b>		<b>Boutonniere deformity</b>	<b>68</b>	<b>38</b>	<b>56%</b>	<b>29</b>	<b>43%</b>	<b>5</b>	<b>11</b>	<b>27</b>	<b>40</b>	<b>47</b>	<b>74</b>	<b>123</b>	<b>179</b>	<b>244</b>	<b>365</b>	<b>47</b>	<b>93.90</b>	<b>32</b>		<b>1</b>	<b>1%</b>
<b>736.29</b>		<b>Other</b>	<b>30</b>	<b>6</b>	<b>20%</b>	<b>22</b>	<b>73%</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>13</b>	<b>26</b>	<b>49</b>	<b>59</b>	<b>191</b>	<b>346</b>	<b>365</b>	<b>26</b>	<b>86.95</b>	<b>26</b>		<b>2</b>	<b>7%</b>
<b>736.6</b>		<b>Other acquired deformities of knee</b>	<b>51</b>	<b>6</b>	<b>12%</b>	<b>28</b>	<b>55%</b>	<b>19</b>	<b>30</b>	<b>60</b>	<b>80</b>	<b>122</b>	<b>137</b>	<b>158</b>	<b>195</b>	<b>295</b>	<b>365</b>	<b>122</b>	<b>128.54</b>	<b>16</b>		<b>17</b>	<b>33%</b>
736.7	I	Other acquired deformities of ankle and foot	46	2	4%	25	54%	20	42	76	95	178	238	291	365	365	365	178	187.08	34		19	41%
<b>736.79</b>		<b>Other</b>	<b>32</b>	<b>1</b>	<b>3%</b>	<b>20</b>	<b>63%</b>	<b>14</b>	<b>31</b>	<b>47</b>	<b>76</b>	<b>100</b>	<b>178</b>	<b>238</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>100</b>	<b>166.60</b>	<b>31</b>		<b>11</b>	<b>34%</b>
736.8	I	Acquired deformities of other parts of limbs	23		0%	16	70%	45	101	151	161	354	365	365	365	365	365	354	249.13	61		7	30%
738	I	Other acquired deformity	105	19	18%	63	60%	10	21	30	50	76	121	216	275	365	365	76	136.57	10	7	23	22%
<b>738.4</b>		<b>Acquired spondylolisthesis</b>	<b>64</b>	<b>10</b>	<b>16%</b>	<b>35</b>	<b>55%</b>	<b>4</b>	<b>12</b>	<b>23</b>	<b>33</b>	<b>50</b>	<b>102</b>	<b>221</b>	<b>269</b>	<b>365</b>	<b>365</b>	<b>50</b>	<b>129.80</b>	<b>30</b>		<b>19</b>	<b>30%</b>
<b>738.8</b>		<b>Acquired deformity of other specified site</b>	<b>33</b>	<b>4</b>	<b>12%</b>	<b>25</b>	<b>76%</b>	<b>17</b>	<b>21</b>	<b>61</b>	<b>69</b>	<b>94</b>	<b>129</b>	<b>206</b>	<b>262</b>	<b>365</b>	<b>365</b>	<b>94</b>	<b>148.76</b>	<b>30</b>		<b>4</b>	<b>12%</b>
739	I	Nonallopathic lesions, not elsewhere classified	515	248	48%	227	44%	1	2	4	6	10	16	28	64	177	365	10	49.09	29		40	8%
<b>739.1</b>		<b>Cervical region</b>	<b>125</b>	<b>74</b>	<b>59%</b>	<b>42</b>	<b>34%</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>8</b>	<b>14</b>	<b>21</b>	<b>78</b>	<b>178</b>	<b>365</b>	<b>8</b>	<b>55.10</b>	<b>8</b>		<b>9</b>	<b>7%</b>
<b>739.2</b>		<b>Thoracic region</b>	<b>54</b>	<b>30</b>	<b>56%</b>	<b>22</b>	<b>41%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>4</b>	<b>5</b>	<b>5</b>	<b>6</b>	<b>26</b>	<b>365</b>	<b>4</b>	<b>26.41</b>	<b>6</b>		<b>2</b>	<b>4%</b>
<b>739.3</b>		<b>Lumbar region</b>	<b>199</b>	<b>80</b>	<b>40%</b>	<b>107</b>	<b>54%</b>	<b>1</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>10</b>	<b>13</b>	<b>23</b>	<b>34</b>	<b>86</b>	<b>365</b>	<b>10</b>	<b>35.27</b>	<b>9</b>		<b>12</b>	<b>6%</b>
<b>739.4</b>		<b>Sacral region</b>	<b>72</b>	<b>32</b>	<b>44%</b>	<b>30</b>	<b>42%</b>	<b>1</b>	<b>2</b>	<b>5</b>	<b>7</b>	<b>20</b>	<b>32</b>	<b>53</b>	<b>73</b>	<b>292</b>	<b>365</b>	<b>20</b>	<b>69.03</b>	<b>16</b>		<b>10</b>	<b>14%</b>
755	I	Other congenital anomalies of limbs	26	3	12%	18	69%	12	30	41	73	78	116	199	286	365	365	78	140.67	16		5	19%
<b>756</b>		<b>Other congenital musculoskeletal anomalies</b>	<b>238</b>	<b>29</b>	<b>12%</b>	<b>108</b>	<b>45%</b>	<b>6</b>															

### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
780	I	General symptoms	121	51	42%	53	44%	1	2	3	4	16	19	40	58	89	365	16	46.60	10	5	17	14%
<b>780.2</b>		<b>Syncope and collapse</b>	<b>69</b>	<b>40</b>	<b>58%</b>	<b>26</b>	<b>38%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>3</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>17</b>	<b>365</b>	<b>3</b>	<b>18.69</b>	<b>9</b>	<b>3</b>	<b>3</b>	<b>4%</b>
782	I	Symptoms involving skin and other integumentary tissue	399	255	64%	118	30%	1	2	3	5	9	14	33	55	178	365	9	47.54	10	6	26	7%
<b>782.0</b>		<b>Disturbance of skin sensation</b>	<b>191</b>	<b>121</b>	<b>63%</b>	<b>57</b>	<b>30%</b>	<b>1</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>14</b>	<b>33</b>	<b>55</b>	<b>98</b>	<b>181</b>	<b>365</b>	<b>14</b>	<b>60.75</b>	<b>10</b>		<b>13</b>	<b>7%</b>
<b>782.1</b>		<b>Rash and other nonspecific skin eruption</b>	<b>119</b>	<b>88</b>	<b>74%</b>	<b>22</b>	<b>18%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>9</b>	<b>1</b>	<b>2.45</b>	<b>2</b>		<b>9</b>	<b>8%</b>
<b>782.2</b>		<b>Localized superficial swelling, mass, or lump</b>	<b>38</b>	<b>18</b>	<b>47%</b>	<b>19</b>	<b>50%</b>	<b>2</b>	<b>4</b>	<b>8</b>	<b>12</b>	<b>20</b>	<b>32</b>	<b>50</b>	<b>61</b>	<b>338</b>	<b>365</b>	<b>20</b>	<b>65.26</b>	<b>9</b>		<b>1</b>	<b>3%</b>
<b>782.3</b>		<b>Edema</b>	<b>45</b>	<b>23</b>	<b>51%</b>	<b>20</b>	<b>44%</b>	<b>1</b>	<b>4</b>	<b>6</b>	<b>10</b>	<b>13</b>	<b>18</b>	<b>28</b>	<b>36</b>	<b>48</b>	<b>365</b>	<b>13</b>	<b>42.65</b>	<b>12</b>	<b>6</b>	<b>2</b>	<b>4%</b>
784	I	Symptoms involving head and neck	219	87	40%	110	50%	2	4	7	12	17	23	52	91	217	365	17	60.89	15	3	22	10%
<b>784.0</b>		<b>Headache</b>	<b>157</b>	<b>54</b>	<b>34%</b>	<b>85</b>	<b>54%</b>	<b>2</b>	<b>6</b>	<b>10</b>	<b>14</b>	<b>21</b>	<b>34</b>	<b>64</b>	<b>99</b>	<b>236</b>	<b>365</b>	<b>21</b>	<b>70.24</b>	<b>2</b>	<b>3</b>	<b>18</b>	<b>11%</b>
<b>784.7</b>		<b>Epistaxis</b>	<b>55</b>	<b>31</b>	<b>56%</b>	<b>22</b>	<b>40%</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>7</b>	<b>12</b>	<b>17</b>	<b>20</b>	<b>42</b>	<b>4</b>	<b>9.00</b>	<b>3</b>		<b>2</b>	<b>4%</b>
785	I	Symptoms involving cardiovascular system	48	10	21%	34	71%	2	10	20	25	51	79	99	220	282	365	51	95.68	7	3	4	8%
<b>785.4</b>		<b>Gangrene</b>	<b>30</b>	<b>1</b>	<b>3%</b>	<b>25</b>	<b>83%</b>	<b>11</b>	<b>20</b>	<b>32</b>	<b>53</b>	<b>79</b>	<b>89</b>	<b>138</b>	<b>220</b>	<b>308</b>	<b>365</b>	<b>79</b>	<b>116.20</b>	<b>42</b>		<b>4</b>	<b>13%</b>
786	I	Symptoms involving respiratory system and other chest symptoms	94	52	55%	35	37%	1	1	2	2	2	3	6	14	31	128	2	12.37	4	3	7	7%
<b>786.0</b>	I	<b>Dyspnea and respiratory abnormalities</b>	<b>51</b>	<b>29</b>	<b>57%</b>	<b>18</b>	<b>35%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>9</b>	<b>16</b>	<b>43</b>	<b>74</b>	<b>3</b>	<b>11.44</b>	<b>8</b>		<b>4</b>	<b>8%</b>
<b>786.5</b>	I	<b>Chest pain</b>	<b>36</b>	<b>18</b>	<b>50%</b>	<b>15</b>	<b>42%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>27</b>	<b>128</b>	<b>2</b>	<b>12.93</b>	<b>9</b>	<b>5</b>	<b>3</b>	<b>8%</b>
800	I	Fracture of vault of skull	46	1	2%	32	70%	6	13	22	33	37	48	72	139	230	365	37	79.22	23		13	28%
800.0	I	Closed without mention of intracranial injury	24		0%	16	67%	6	10	13	20	23	34	37	72	186	262	23	53.75	23		8	33%
801	I	Fracture of base of skull	88	15	17%	60	68%	5	7	13	23	29	33	51	104	173	357	29	59.55	26		13	15%
801.0	I	Closed without mention of intracranial injury	53	10	19%	39	74%	4	5	8	11	22	27	31	54	118	263	22	38.79	25		4	8%
<b>801.01</b>		<b>Closed without mention of intracranial injury</b>	<b>29</b>	<b>7</b>	<b>24%</b>	<b>22</b>	<b>76%</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>8</b>	<b>11</b>	<b>22</b>	<b>25</b>	<b>28</b>	<b>37</b>	<b>263</b>	<b>11</b>	<b>27.77</b>	<b>25</b>			<b>0%</b>
802	I	Fracture of face bones	1,291	538	42%	676	52%	1	2	3	4	6	11	16	24	44	365	6	18.98	11		77	6%
<b>802.0</b>		<b>Nasal bones, closed</b>	<b>860</b>	<b>421</b>	<b>49%</b>	<b>393</b>	<b>46%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>12</b>	<b>21</b>	<b>365</b>	<b>3</b>	<b>10.77</b>	<b>4</b>		<b>46</b>	<b>5%</b>
<b>802.1</b>		<b>Nasal bones, open</b>	<b>52</b>	<b>27</b>	<b>52%</b>	<b>22</b>	<b>42%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>14</b>	<b>15</b>	<b>21</b>	<b>89</b>	<b>4</b>	<b>11.55</b>	<b>7</b>		<b>3</b>	<b>6%</b>
802.2	I	Mandible, closed	85	20	24%	58	68%	2	6	12	15	20	24	35	44	54	263	20	29.40	20		7	8%
<b>802.20</b>		<b>Unspecified site</b>	<b>34</b>	<b>8</b>	<b>24%</b>	<b>24</b>	<b>71%</b>	<b>2</b>	<b>7</b>	<b>13</b>	<b>15</b>	<b>18</b>	<b>23</b>	<b>24</b>	<b>40</b>	<b>54</b>	<b>263</b>	<b>18</b>	<b>31.83</b>	<b>18</b>		<b>2</b>	<b>6%</b>
<b>802.4</b>		<b>Malar and maxillary bones, closed</b>	<b>107</b>	<b>31</b>	<b>29%</b>	<b>69</b>	<b>64%</b>	<b>1</b>	<b>2</b>	<b>5</b>	<b>9</b>	<b>17</b>	<b>19</b>	<b>25</b>	<b>38</b>	<b>69</b>	<b>365</b>	<b>17</b>	<b>28.70</b>	<b>17</b>		<b>7</b>	<b>7%</b>
<b>802.6</b>		<b>Orbital floor (blow-out), closed</b>	<b>48</b>	<b>12</b>	<b>25%</b>	<b>35</b>	<b>73%</b>	<b>3</b>	<b>6</b>	<b>7</b>	<b>9</b>	<b>13</b>	<b>15</b>	<b>25</b>	<b>27</b>	<b>47</b>	<b>148</b>	<b>13</b>	<b>21.40</b>	<b>16</b>		<b>1</b>	<b>2%</b>
<b>802.8</b>		<b>Other facial bones, closed</b>	<b>101</b>	<b>21</b>	<b>21%</b>	<b>70</b>	<b>69%</b>	<b>3</b>	<b>6</b>	<b>9</b>	<b>13</b>	<b>16</b>	<b>21</b>	<b>28</b>	<b>36</b>	<b>58</b>	<b>186</b>	<b>16</b>	<b>27.60</b>	<b>16</b>		<b>10</b>	<b>10%</b>
805	I	Fracture of vertebral column without mention of spinal cord injury	1,320	221	17%	906	69%	3	6	17	31	48	72	106	159	284	365	48	91.73	42		193	15%
805.0	I	Cervical, closed	122	13	11%	76	62%	4	20	45	63	92	132	199	280	365	365	92	140.11	36		33	27%
<b>805.07</b>		<b>Cervical, closed</b>	<b>42</b>	<b>7</b>	<b>17%</b>	<b>25</b>	<b>60%</b>	<b>3</b>	<b>5</b>	<b>46</b>	<b>61</b>	<b>69</b>	<b>136</b>	<b>199</b>	<b>305</b>	<b>365</b>	<b>365</b>	<b>69</b>	<b>141.36</b>	<b>36</b>		<b>10</b>	<b>24%</b>
<b>805.2</b>		<b>Dorsal [thoracic], closed</b>	<b>332</b>	<b>46</b>	<b>14%</b>	<b>230</b>	<b>69%</b>	<b>6</b>	<b>19</b>	<b>31</b>	<b>48</b>	<b>71</b>	<b>100</b>	<b>131</b>	<b>195</b>	<b>365</b>	<b>365</b>	<b>71</b>	<b>114.38</b>	<b>59</b>		<b>56</b>	<b>17%</b>
<b>805.4</b>		<b>Lumbar, closed</b>	<b>516</b>	<b>60</b>	<b>12%</b>	<b>370</b>	<b>72%</b>	<b>5</b>	<b>11</b>	<b>24</b>	<b>43</b>	<b>61</b>	<b>93</b>	<b>123</b>	<b>177</b>	<b>250</b>	<b>365</b>	<b>61</b>	<b>99.39</b>	<b>60</b>		<b>86</b>	<b>17%</b>
<b>805.6</b>		<b>Sacrum and coccyx, closed</b>	<b>345</b>	<b>102</b>	<b>30%</b>	<b>228</b>	<b>66%</b>	<b>1</b>	<b>3</b>	<b>4</b>	<b>7</b>	<b>13</b>	<b>25</b>	<b>38</b>	<b>50</b>	<b>84</b>	<b>365</b>	<b>13</b>	<b>39.49</b>	<b>22</b>		<b>15</b>	<b>4%</b>
807	I	Fracture of rib(s), sternum, larynx, and trachea	1,928	636	33%	1,207	63%	2	3	4	7	12	18	27	39	58	365	12	27.23	16		85	4%
807.0	I	Rib(s), closed	1,846	615	33%	1,150	62%	2	3	4	7	12	18	26	38	56	365	12	26.47	13		81	4%
<b>807.00</b>		<b>Rib(s), closed</b>	<b>330</b>	<b>135</b>	<b>41%</b>	<b>188</b>	<b>57%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>7</b>	<b>13</b>	<b>20</b>	<b>34</b>	<b>116</b>	<b>6</b>	<b>12.94</b>	<b>13</b>		<b>7</b>	<b>2%</b>
<b>807.01</b>		<b>Rib(s), closed</b>	<b>914</b>	<b>336</b>	<b>37%</b>	<b>533</b>	<b>58%</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>11</b>	<b>16</b>	<b>22</b>	<b>33</b>	<b>53</b>	<b>365</b>	<b>11</b>	<b>24.93</b>	<b>13</b>		<b>45</b>	<b>5%</b>
<b>807.02</b>		<b>Rib(s), closed</b>	<b>318</b>	<b>84</b>	<b>26%</b>	<b>222</b>	<b>70%</b>	<b>2</b>	<b>4</b>	<b>5</b>	<b>10</b>	<b>14</b>	<b>22</b>	<b>30</b>	<b>39</b>	<b>56</b>	<b>365</b>	<b>14</b>	<b>27.04</b>	<b>13</b>		<b>12</b>	<b>4%</b>
<b>807.03</b>		<b>Rib(s), closed</b>	<b>154</b>	<b>36</b>	<b>23%</b>	<b>115</b>	<b>75%</b>	<b>2</b>	<b>6</b>	<b>11</b>	<b>14</b>	<b>20</b>	<b>26</b>	<b>36</b>	<b>49</b>	<b>71</b>	<b>365</b>	<b>20</b>	<b>33.68</b>	<b>13</b>		<b>3</b>	<b>2%</b>
<b>807.04</b>		<b>Rib(s), closed</b>	<b>45</b>	<b>10</b>	<b>22%</b>	<b>32</b>	<b>71%</b>	<b>12</b>	<b>19</b>	<b>30</b>	<b>34</b>	<b>44</b>	<b>51</b>	<b>61</b>	<b>99</b>	<b>182</b>	<b>365</b>	<b>44</b>	<b>73.75</b>	<b>13</b>		<b>3</b>	<b>7%</b>
<b>807.09</b>		<b>Rib(s), closed</b>	<b>38</b>	<b>5</b>	<b>13%</b>	<b>28</b>	<b>74%</b>	<b>7</b>	<b>17</b>	<b>32</b>	<b>38</b>	<b>41</b>	<b>50</b>	<b>62</b>	<b>100</b>	<b>147</b>	<b>189</b>	<b>41</b>	<b>60.61</b>	<b>13</b>		<b>5</b>	<b>13%</b>
<b>807.2</b>		<b>Sternum, closed</b>	<b>57</b>	<b>16</b>	<b>28%</b>	<b>38</b>	<b>67%</b>	<b>1</b>	<b>3</b>	<b>5</b>	<b>6</b>	<b>17</b>	<b>34</b>	<b>57</b>	<b>77</b>	<b>111</b>	<b>354</b>	<b>17</b>	<b>45.08</b>	<b>10</b>		<b>3</b>	<b>5%</b>
808	I	Fracture of pelvis	394	27	7%	294	75%	17	32	52	65	85	108	135	184	365	365	85	121.04	63		73	19%
<b>808.0</b>		<b>Acetabulum, closed</b>	<b>150</b>	<b>7</b>	<b>5%</b>	<b>113</b>	<b>75%</b>	<b>24</b>	<b>55</b>	<b>80</b>	<b>96</b>	<b>117</b>	<b>137</b>	<b>180</b>	<b>277</b>	<b>365</b>	<b>365</b>	<b>117</b>	<b>151.58</b>	<b>93</b>		<b>30</b>	<b>20%</b>
<b>808.2</b>		<b>Pubis, closed</b>	<b>152</b>	<b>15</b>	<b>10%</b>	<b>115</b>	<b>76%</b>	<b>7</b>	<b>27</b>	<b>43</b>	<b>53</b>	<b>63</b>	<b>81</b>	<b>107</b>	<b>141</b>	<b>222</b>	<b>365</b>	<b>63</b>	<b>93.83</b>	<b>67</b>		<b>22</b>	<b>14%</b>
808.4	I	Other specified part, closed	68	5	7%	48	71%	19	29	38	52	62	93	115	183</								

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
810.02		Closed	62	6	10%	49	79%	4	12	20	33	53	59	73	140	186	357	53	75.47	26		7	11%
810.03		Closed	48	11	23%	34	71%	3	13	20	29	38	39	42	73	139	256	38	51.68	26		3	6%
811	I	Fracture of scapula	193	35	18%	137	71%	4	6	19	31	44	61	74	94	213	365	44	75.93	31	15	21	11%
811.0	I	Closed	193	35	18%	137	71%	4	6	19	31	44	61	74	94	213	365	44	75.93	29		21	11%
811.00		Closed	90	13	14%	67	74%	3	5	14	29	41	57	64	80	152	365	41	60.93	29		10	11%
811.03		Closed	65	15	23%	43	66%	6	15	20	25	49	74	94	174	345	365	49	96.93	29		7	11%
811.09		Closed	18		0%	16	89%	5	11	36	45	47	71	79	114	365	365	47	94.75	29		2	11%
812	I	Fracture of humerus	1,231	234	19%	887	72%	4	9	23	38	55	72	91	128	213	365	55	82.33	18		110	9%
812.0		Upper end, closed	639	103	16%	491	77%	4	12	26	46	60	76	95	132	217	365	60	85.70	24		45	7%
812.00		Upper end, unspecified part	163	31	19%	124	76%	4	6	14	32	42	60	73	96	141	365	42	61.89	18		8	5%
812.01		Surgical neck	128	13	10%	104	81%	5	23	45	55	62	77	89	122	225	365	62	93.26	31		11	9%
812.03		Greater tuberosity	206	35	17%	158	77%	5	16	26	49	72	91	127	160	272	365	72	98.82	19		13	6%
812.09		Other	125	22	18%	91	73%	3	7	30	50	62	76	95	131	228	365	62	88.14	20		12	10%
812.2	I	Shaft or unspecified part, closed	241	37	15%	175	73%	4	11	26	49	66	86	110	153	215	365	66	92.82	20		29	12%
812.20		Unspecified part of humerus	127	27	21%	85	67%	3	6	13	28	54	68	103	150	188	365	54	78.73	13		15	12%
812.21		Shaft of humerus	114	10	9%	90	79%	10	24	41	62	76	94	120	157	252	365	76	106.12	26		14	12%
812.4	I	Lower end, closed	306	92	30%	190	62%	2	5	9	21	31	47	59	81	125	365	31	52.65	16		24	8%
812.40		Lower end, unspecified part	199	70	35%	112	56%	2	4	6	15	23	40	48	68	109	365	23	44.38	17		17	9%
812.41		Supracondylar fracture of humerus	21	4	19%	16	76%	6	7	11	25	32	60	73	99	334	365	32	80.75	22		1	5%
812.42		Lateral condyle	33	7	21%	23	70%	2	6	8	20	31	42	64	93	144	202	31	51.13	20		3	9%
812.43		Medial condyle	25	5	20%	19	76%	1	11	21	31	31	53	58	70	127	197	31	50.21	23		1	4%
812.5	I	Lower end, open	27		0%	22	81%	4	8	13	24	40	74	122	334	365	365	40	117.91	22		5	19%
813	I	Fracture of radius and ulna	5,187	1,323	26%	3,561	69%	2	5	10	19	32	47	62	87	130	365	32	55.14	29		303	6%
813.0	I	Upper end, closed	1,769	497	28%	1,171	66%	2	4	6	13	23	35	47	62	96	365	23	42.21	24		101	6%
813.01		Olecranon process of ulna	285	76	27%	185	65%	2	5	11	20	35	52	73	101	153	365	35	58.86	33		24	8%
813.02		Coronoid process of ulna	54	15	28%	35	65%	1	4	15	15	22	36	64	81	106	319	22	48.29	25		4	7%
813.04		Other and unspecified fractures of proximal end of ulna (alone)	48	18	38%	28	58%	3	3	6	38	45	60	84	113	159	365	45	73.00	45		2	4%
813.05		Head of radius	1,119	314	28%	744	66%	2	3	6	11	19	29	42	55	80	365	19	34.99	21		61	5%
813.06		Neck of radius	129	35	27%	89	69%	2	4	13	20	27	34	45	55	63	363	27	39.96	27		5	4%
813.07		Other and unspecified fractures of proximal end of radius (alone)	80	26	33%	52	65%	2	2	4	5	8	28	44	56	66	214	8	29.13	10		2	3%
813.1	I	Upper end, open	37	11	30%	22	59%	6	16	28	50	77	95	106	119	192	365	77	95.55	81		4	11%
813.2	I	Shaft, closed	150	28	19%	109	73%	3	6	13	21	33	50	61	102	132	365	33	55.58	33		13	9%
813.21		Radius (alone)	64	12	19%	46	72%	4	6	14	31	39	55	67	96	132	230	39	55.59	46		6	9%
813.22		Ulna (alone)	48	11	23%	32	67%	1	5	6	15	19	41	48	59	71	365	19	41.63	21		5	10%
813.23		Radius with ulna	36	5	14%	30	83%	3	11	15	28	31	76	102	113	184	307	31	72.17	28		1	3%
813.3	I	Shaft, open	34	2	6%	23	68%	32	61	62	115	131	179	289	365	365	365	131	177.30	78		9	26%
813.4	I	Lower end, closed	2,817	662	24%	2,002	71%	3	6	12	24	40	55	72	95	138	365	40	60.77	39		153	5%
813.41		Colles' fracture	394	74	19%	302	77%	3	8	15	24	43	62	77	96	151	365	43	62.51	39		18	5%
813.42		Other fractures of distal end of radius (alone)	1,886	444	24%	1,338	71%	2	5	11	24	41	55	71	95	142	365	41	61.17	39		104	6%
813.43		Distal end of ulna (alone)	217	90	41%	118	54%	3	4	6	9	17	28	50	68	101	328	17	37.95	31		9	4%
813.44		Radius with ulna, lower end	302	48	16%	232	77%	4	9	20	31	46	61	86	115	139	365	46	68.61	45		22	7%
813.5	I	Lower end, open	49	3	6%	39	80%	6	12	18	31	54	82	91	148	179	365	54	83.28	53		7	14%
813.52		Other fractures of distal end of radius (alone)	25	1	4%	20	80%	6	12	15	18	45	79	104	148	154	365	45	82.00	45		4	16%
813.8	I	Unspecified part, closed	317	118	37%	184	58%	2	3	5	10	18	40	61	81	131	365	18	49.45	19		15	5%
813.81		Radius (alone)	191	75	39%	106	55%	2	3	5	8	19	35	53	72	118	291	19	43.74	18		10	5%
813.82		Ulna (alone)	55	18	33%	35	64%	1	2	4	4	7	10	15	43	99	365	7	38.34	13		2	4%
813.83		Radius with ulna	49	14	29%	34	69%	6	11	33	58	65	92	105	136	164	365	65	81.71	65		1	2%
814	I	Fracture of carpal bone(s)	1,486	610	41%	806	54%	2	3	5	10	19	37	55	78	123	365	19	49.15	18		70	5%
814.0	I	Closed	1,465	602	41%	794	54%	2	3	5	9	19	36	54	78	120	365	19	48.01	16		69	5%
814.00		Closed	493	225	46%	250	51%	1	2	3	5	6	14	32	56	86	338	6	32.24	16		18	4%
814.01		Closed	630	242	38%	353	56%	2	4	8	16	30	51	69	97	135	365	30	58.64	16		35	6%

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
814.02		Closed	46	17	37%	24	52%	1	2	4	6	12	60	70	193	279	365	12	87.38	16		5	11%
814.03		Closed	162	70	43%	90	56%	1	3	6	12	25	33	45	61	81	350	25	38.38	16		2	1%
814.08		Closed	35	11	31%	22	63%	2	3	8	20	29	39	44	59	90	239	29	40.82	16		2	6%
814.09		Closed	24	7	29%	15	63%	3	3	24	31	46	48	59	83	142	256	46	58.60	16		2	8%
815	I	Fracture of metacarpal bone(s)	2,426	929	38%	1,374	57%	2	3	5	11	19	32	44	57	87	365	19	37.19	21		123	5%
815.0	I	Closed	2,293	901	39%	1,281	56%	2	3	5	10	18	31	42	55	79	365	18	33.80	20		111	5%
815.00		Closed	1,297	541	42%	703	54%	2	3	5	7	14	25	38	50	73	365	14	29.91	20		53	4%
815.01		Closed	111	40	36%	63	57%	1	3	4	10	18	30	47	67	101	267	18	39.21	20		8	7%
815.02		Closed	296	116	39%	167	56%	2	3	6	14	23	36	43	57	75	319	23	34.86	20		13	4%
815.03		Closed	318	114	36%	180	57%	2	4	8	14	23	38	48	56	91	250	23	38.37	20		24	8%
815.04		Closed	221	74	33%	134	61%	3	6	12	25	33	44	52	66	96	230	33	42.29	20		13	6%
815.09		Closed	50	16	32%	34	68%	2	2	3	4	6	15	48	65	118	365	6	41.38	20			0%
815.1	I	Open	133	28	21%	93	70%	3	5	15	30	33	55	89	143	286	365	33	83.88	28		12	9%
815.10		Open	52	15	29%	32	62%	3	4	5	12	28	32	36	80	147	290	28	46.72	28		5	10%
815.13		Open	40	6	15%	28	70%	5	24	32	38	43	100	143	248	365	365	43	117.11	28		6	15%
816	I	Fracture of one or more phalanges of hand	13,652	7,493	55%	5,762	42%	1	2	3	4	7	13	21	34	60	365	7	23.65	13		397	3%
816.0	I	Closed	9,925	5,995	60%	3,652	37%	1	2	2	3	5	9	17	28	51	365	5	18.98	6		278	3%
816.00		Closed	4,163	2,699	65%	1,350	32%	1	1	2	3	4	6	10	20	40	365	4	14.80	6		114	3%
816.01		Middle or proximal phalanx or phalanges	1,972	1,026	52%	888	45%	2	2	4	6	12	20	32	49	77	365	12	29.43	13		58	3%
816.02		Distal phalanx or phalanges	3,709	2,232	60%	1,371	37%	1	2	2	3	5	7	13	24	40	365	5	15.85	6		106	3%
816.03		Distal phalanx or phalanges	81	38	47%	43	53%	1	2	6	7	10	18	33	55	75	223	10	34.30	6			0%
816.1	I	Open	3,727	1,498	40%	2,110	57%	1	3	4	8	13	20	31	48	76	365	13	31.74	13		119	3%
816.10		Open	814	397	49%	395	49%	1	2	3	4	7	12	20	33	61	365	7	23.23	13		22	3%
816.11		Open	632	132	21%	473	75%	2	6	12	20	31	48	62	85	146	365	31	57.89	13		27	4%
816.12		Open	2,214	950	43%	1,199	54%	1	3	4	6	11	17	25	36	53	365	11	22.22	13		65	3%
816.13		Open	67	19	28%	43	64%	2	5	17	30	41	62	101	146	264	365	41	87.77	13		5	7%
817	I	Multiple fractures of hand bones	88	51	58%	35	40%	2	4	4	6	16	28	42	47	71	92	16	27.20	21		2	2%
817.0		Closed	64	39	61%	24	38%	2	4	5	7	16	41	47	63	76	92	16	31.21	33		1	2%
820	I	Fracture of neck of femur	466	16	3%	349	75%	39	61	76	88	103	126	161	217	319	365	103	138.64	45		101	22%
820.0	I	Transcervical fracture, closed	89	3	3%	70	79%	38	53	73	79	87	95	114	144	223	365	87	112.23	88		16	18%
820.09		Other	48	3	6%	37	77%	31	60	76	79	88	94	102	124	223	291	88	102.32	86		8	17%
820.2	I	Pertrochanteric fracture, closed	194	6	3%	143	74%	43	68	82	98	117	132	162	215	313	365	117	142.48	105		45	23%
820.20		Trochanteric section, unspecified	24	2	8%	19	79%	8	20	41	51	81	93	118	122	131	139	81	73.63	81		3	13%
820.21		Intertrochanteric section	140	2	1%	106	76%	52	72	85	105	120	141	177	228	354	365	120	153.43	105		32	23%
820.22		Subtrochanteric section	30	2	7%	18	60%	58	72	75	107	145	165	193	210	263	365	145	150.61	118		10	33%
820.8		Unspecified part of neck of femur, closed	173	7	4%	128	74%	35	62	76	88	108	132	177	260	342	365	108	146.77	92		38	22%
821	I	Fracture of other and unspecified parts of femur	334	30	9%	229	69%	13	40	63	78	104	141	180	237	365	365	104	140.13	53		75	22%
821.0	I	Shaft or unspecified part, closed	121	10	8%	83	69%	10	57	71	86	107	132	164	211	305	365	107	134.61	88		28	23%
821.00		Unspecified part of femur	44	7	16%	31	70%	6	10	57	76	81	93	129	152	200	365	81	101.35	81		6	14%
821.01		Shaft	77	3	4%	52	68%	42	64	78	104	129	141	193	268	340	365	129	154.44	104		22	29%
821.1	I	Shaft or unspecified part, open	23		0%	18	78%	23	107	125	188	191	228	270	315	365	365	191	198.44	125		5	22%
821.2	I	Lower end, closed	171	19	11%	117	68%	11	34	51	68	83	113	165	220	365	365	83	127.37	76		35	20%
821.20		Lower end, unspecified part	30	4	13%	20	67%	2	37	67	76	113	145	217	236	333	365	113	150.15	107		6	20%
821.21		Condyle, femoral	116	13	11%	84	72%	16	33	49	62	77	96	164	219	365	365	77	124.38	66		19	16%
822	I	Fracture of patella	720	117	16%	545	76%	4	9	21	36	51	66	82	112	179	365	51	73.30	37		58	8%
822.0		Closed	708	115	16%	536	76%	4	10	21	36	51	66	82	113	180	365	51	73.49	39		57	8%
823	I	Fracture of tibia and fibula	1,806	229	13%	1,391	77%	6	20	36	55	75	97	124	173	270	365	75	105.92	37		186	10%
823.0	I	Upper end, closed	662	74	11%	518	78%	8	23	42	55	72	89	117	153	250	365	72	100.78	40		70	11%
823.00		Upper end, closed	470	47	10%	363	77%	9	34	51	67	80	98	124	163	267	365	80	110.21	40		60	13%
823.01		Upper end, closed	156	24	15%	127	81%	3	11	22	32	42	53	68	97	174	365	42	64.87	40		5	3%
823.02		Upper end, closed	36	3	8%	28	78%	14	26	74	106	122	159	186	219	365	365	122	141.32	40		5	14%

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
823.1	I	Upper end, open	30	4	13%	19	63%	3	59	87	115	152	213	240	253	365	365	152	163.84	37		7	23%
823.2	I	Shaft, closed	341	45	13%	256	75%	7	25	42	61	92	109	132	180	283	365	92	112.78	41		40	12%
<b>823.20</b>		<b>Shaft, closed</b>	<b>96</b>	<b>15</b>	<b>16%</b>	<b>69</b>	<b>72%</b>	<b>8</b>	<b>36</b>	<b>52</b>	<b>78</b>	<b>96</b>	<b>116</b>	<b>146</b>	<b>201</b>	<b>365</b>	<b>365</b>	<b>96</b>	<b>124.45</b>	<b>41</b>		<b>12</b>	<b>13%</b>
823.21		Shaft, closed	122	24	20%	88	72%	4	8	20	32	41	51	65	92	114	226	41	51.92	41		10	8%
823.22		Shaft, closed	123	6	5%	99	80%	26	64	95	110	127	163	191	275	341	365	127	158.75	41		18	15%
823.3	I	Shaft, open	80	2	3%	64	80%	54	104	135	170	207	259	320	365	365	365	207	217.52	54		14	18%
823.32		Shaft, open	57	1	2%	46	81%	73	135	150	195	245	277	340	365	365	365	245	235.17	54		10	18%
823.8	I	Unspecified part, closed	636	102	16%	494	78%	4	9	22	39	58	77	101	141	227	365	58	87.10	21		40	6%
823.80		Unspecified part, closed	188	33	18%	143	76%	3	7	20	44	56	76	103	130	206	365	56	85.29	21		12	6%
823.81		Unspecified part, closed	337	64	19%	253	75%	3	6	17	26	43	59	76	99	156	365	43	65.36	21		20	6%
823.82		Unspecified part, closed	111	5	5%	98	88%	16	34	76	105	126	153	181	234	334	365	126	145.85	21		8	7%
823.9	I	Unspecified part, open	57	2	4%	40	70%	20	32	55	96	131	163	204	256	365	365	131	155.05	45		15	26%
823.92		Unspecified part, open	41	1	2%	27	66%	51	96	131	138	172	246	247	365	365	365	172	197.67	45		13	32%
824	I	Fracture of ankle	3,309	510	15%	2,601	79%	4	10	20	34	49	64	82	103	153	365	49	68.07	35		198	6%
824.0		Medial malleolus, closed	471	62	13%	380	81%	5	16	29	44	56	71	83	104	165	365	56	74.69	52		29	6%
824.2		Lateral malleolus, closed	1,040	164	16%	837	80%	3	9	16	27	39	52	66	82	114	365	39	52.51	22		39	4%
824.4		Bimalleolar, closed	340	10	3%	302	89%	20	42	56	69	82	95	108	134	191	365	82	96.92	76		28	8%
824.5		Bimalleolar, open	23	1	4%	16	70%	70	109	115	117	138	191	238	244	365	365	138	183.56	109		6	26%
824.6		Trimalleolar, closed	203	8	4%	172	85%	19	38	62	76	89	106	128	157	195	365	89	108.08	73		23	11%
824.8		Unspecified, closed	1,178	258	22%	855	73%	2	5	9	18	33	48	65	92	145	365	33	55.59	29		65	6%
824.9		Unspecified, open	31	2	6%	23	74%	61	89	110	207	281	297	333	354	365	365	281	224.43	67		6	19%
825	I	Fracture of one or more tarsal and metatarsal bones	4,483	1,027	23%	3,201	71%	2	5	11	21	35	49	66	91	143	365	35	59.00	34		255	6%
825.0		Fracture of calcaneus, closed	636	75	12%	493	78%	6	25	46	67	86	108	137	184	295	365	86	114.74	59		68	11%
825.2	I	Fracture of other tarsal and metatarsal bones, closed	3,765	940	25%	2,650	70%	2	4	8	17	29	42	55	74	110	365	29	47.61	30		175	5%
825.20		Unspecified bone(s) of foot [except toes]	384	143	37%	218	57%	1	2	3	6	10	20	34	54	90	365	10	34.01	10		23	6%
825.21		Astragalus	308	69	22%	223	72%	2	5	9	19	27	37	52	85	132	365	27	51.87	25		16	5%
825.22		Navicular [scaphoid], foot	192	41	21%	141	73%	2	4	11	18	31	43	53	75	119	365	31	49.70	30		10	5%
825.23		Cuboid	165	39	24%	118	72%	3	4	7	14	26	36	45	62	102	365	26	41.21	29		8	5%
825.24		Cuneiform, foot	77	11	14%	63	82%	5	8	15	30	50	62	81	97	138	365	50	63.00	48		3	4%
825.25		Metatarsal bone(s)	2,558	615	24%	1,835	72%	2	4	9	18	31	45	56	74	107	365	31	47.42	32		108	4%
825.29		Other	81	22	27%	52	64%	5	13	19	38	53	68	111	153	207	365	53	83.25	48		7	9%
825.3	I	Fracture of other tarsal and metatarsal bones, open	67	11	16%	50	75%	5	15	25	46	68	85	101	140	242	365	68	90.44	54		6	9%
825.35		Metatarsal bone(s)	35	5	14%	30	86%	20	25	33	54	72	86	95	146	261	365	72	103.37	72			0%
826	I	Fracture of one or more phalanges of foot	3,278	1,446	44%	1,735	53%	1	2	3	5	7	12	20	30	48	365	7	19.27	12		97	3%
826.0		Closed	3,024	1,388	46%	1,548	51%	1	2	3	5	6	11	18	27	45	365	6	18.17	8		88	3%
826.1		Open	254	58	23%	187	74%	2	3	6	12	21	29	36	49	69	159	21	28.44	21		9	4%
831	I	Dislocation of shoulder	1,238	376	30%	788	64%	2	3	6	11	20	32	51	86	168	365	20	55.15	15	28	74	6%
831.0	I	Closed dislocation	1,211	364	30%	774	64%	2	3	6	11	20	33	51	89	170	365	20	55.70	11		73	6%
831.00		Closed dislocation	448	156	35%	270	60%	1	2	4	6	11	20	39	61	135	365	11	43.98	11		22	5%
831.01		Closed dislocation	287	69	24%	194	68%	2	4	6	12	21	37	59	97	173	365	21	58.61	11		24	8%
831.04		Acromioclavicular joint dislocation	433	132	30%	277	64%	2	4	8	15	26	37	54	100	181	365	26	62.88	16		24	6%
832	I	Dislocation of elbow	143	14	10%	122	85%	4	13	23	31	40	50	76	115	157	365	40	73.21	23		7	5%
832.0	I	Closed dislocation	138	14	10%	117	85%	4	10	23	30	38	49	72	102	156	365	38	68.32	27		7	5%
832.00		Closed dislocation	61	8	13%	51	84%	3	6	21	26	30	36	48	66	94	247	30	44.86	27		2	3%
832.02		Closed dislocation	51	3	6%	44	86%	4	14	27	38	45	59	76	107	143	365	45	67.11	27		4	8%
833	I	Dislocation of wrist	142	30	21%	91	64%	5	22	33	63	95	133	181	293	365	365	95	140.22	27		21	15%
833.0	I	Closed dislocation	128	29	23%	84	66%	5	19	30	61	89	129	159	222	365	365	89	127.36	24		15	12%
833.00		Closed dislocation	29	11	38%	17	59%	4	5	19	29	91	117	122	136	173	215	91	81.47	24		1	3%
833.01		Closed dislocation	26	5	19%	16	62%	24	63	77	95	116	172	232	365	365	365	116	167.06	24		5	19%
833.04		Closed dislocation	26	6	23%	16	62%	30	41	46	68	89	132	293	365	365	365	89	162.81	24		4	15%
834	I	Dislocation of finger	750	465	62%	260	35%	1	2	3	4	7	13	23	37	62	365	7	24.69	6		25	3%



**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
834.0	I	Closed dislocation	674	433	64%	223	33%	1	2	3	4	6	10	18	31	52	365	6	20.09	5		18	3%
<b>834.00</b>		<b>Closed dislocation</b>	<b>355</b>	<b>234</b>	<b>66%</b>	<b>109</b>	<b>31%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>11</b>	<b>19</b>	<b>35</b>	<b>365</b>	<b>4</b>	<b>15.43</b>	<b>5</b>		<b>12</b>	<b>3%</b>
<b>834.01</b>		<b>Closed dislocation</b>	<b>94</b>	<b>50</b>	<b>53%</b>	<b>42</b>	<b>45%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>10</b>	<b>14</b>	<b>23</b>	<b>38</b>	<b>66</b>	<b>364</b>	<b>10</b>	<b>28.45</b>	<b>5</b>		<b>2</b>	<b>2%</b>
<b>834.02</b>		<b>Closed dislocation</b>	<b>225</b>	<b>149</b>	<b>66%</b>	<b>72</b>	<b>32%</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>8</b>	<b>16</b>	<b>25</b>	<b>40</b>	<b>55</b>	<b>161</b>	<b>8</b>	<b>22.26</b>	<b>5</b>		<b>4</b>	<b>2%</b>
834.1	I	Open dislocation	76	32	42%	37	49%	4	6	14	19	28	53	62	81	126	250	28	52.41	22		7	9%
<b>834.12</b>		<b>Open dislocation</b>	<b>35</b>	<b>10</b>	<b>29%</b>	<b>22</b>	<b>63%</b>	<b>6</b>	<b>12</b>	<b>14</b>	<b>19</b>	<b>24</b>	<b>48</b>	<b>55</b>	<b>72</b>	<b>199</b>	<b>250</b>	<b>24</b>	<b>56.50</b>	<b>22</b>		<b>3</b>	<b>9%</b>
835	I	Dislocation of hip	21	3	14%	15	71%	11	12	27	48	66	81	87	104	180	365	66	86.27	21		3	14%
835.0	I	Closed dislocation	21	3	14%	15	71%	11	12	27	48	66	81	87	104	180	365	66	86.27	27		3	14%
836	I	Dislocation of knee	5,785	963	17%	4,291	74%	6	18	29	41	55	73	97	134	207	365	55	85.51	26	27	531	9%
<b>836.0</b>		<b>Tear of medial cartilage or meniscus of knee, current</b>	<b>3,542</b>	<b>623</b>	<b>18%</b>	<b>2,644</b>	<b>75%</b>	<b>7</b>	<b>18</b>	<b>29</b>	<b>39</b>	<b>53</b>	<b>68</b>	<b>91</b>	<b>124</b>	<b>182</b>	<b>365</b>	<b>53</b>	<b>79.27</b>	<b>17</b>		<b>275</b>	<b>8%</b>
<b>836.1</b>		<b>Tear of lateral cartilage or meniscus of knee, current</b>	<b>1,587</b>	<b>184</b>	<b>12%</b>	<b>1,209</b>	<b>76%</b>	<b>11</b>	<b>25</b>	<b>39</b>	<b>55</b>	<b>74</b>	<b>96</b>	<b>126</b>	<b>172</b>	<b>270</b>	<b>365</b>	<b>74</b>	<b>106.74</b>	<b>26</b>		<b>194</b>	<b>12%</b>
<b>836.2</b>		<b>Other tear of cartilage or meniscus of knee, current</b>	<b>241</b>	<b>49</b>	<b>20%</b>	<b>164</b>	<b>68%</b>	<b>4</b>	<b>13</b>	<b>23</b>	<b>36</b>	<b>49</b>	<b>62</b>	<b>86</b>	<b>128</b>	<b>244</b>	<b>365</b>	<b>49</b>	<b>85.70</b>	<b>31</b>		<b>28</b>	<b>12%</b>
<b>836.3</b>		<b>Dislocation of patella, closed</b>	<b>357</b>	<b>89</b>	<b>25%</b>	<b>243</b>	<b>68%</b>	<b>2</b>	<b>4</b>	<b>7</b>	<b>15</b>	<b>28</b>	<b>37</b>	<b>55</b>	<b>78</b>	<b>149</b>	<b>365</b>	<b>28</b>	<b>54.82</b>	<b>16</b>		<b>25</b>	<b>7%</b>
836.5	I	Other dislocation of knee, closed	50	14	28%	28	56%	1	2	4	5	7	11	32	44	60	244	7	31.07	7		8	16%
<b>836.50</b>		<b>Dislocation of knee, unspecified</b>	<b>46</b>	<b>13</b>	<b>28%</b>	<b>25</b>	<b>54%</b>	<b>1</b>	<b>1</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>9</b>	<b>14</b>	<b>32</b>	<b>52</b>	<b>60</b>	<b>6</b>	<b>15.68</b>	<b>7</b>		<b>8</b>	<b>17%</b>
837	I	Dislocation of ankle	138	5	4%	106	77%	17	43	65	82	94	117	146	192	293	365	94	124.87	67		27	20%
<b>837.0</b>		<b>Closed dislocation</b>	<b>133</b>	<b>5</b>	<b>4%</b>	<b>104</b>	<b>78%</b>	<b>17</b>	<b>43</b>	<b>66</b>	<b>82</b>	<b>94</b>	<b>117</b>	<b>144</b>	<b>193</b>	<b>293</b>	<b>365</b>	<b>94</b>	<b>125.34</b>	<b>80</b>		<b>24</b>	<b>18%</b>
838	I	Dislocation of foot	143	28	20%	103	72%	8	25	43	60	92	124	156	201	309	365	92	120.50	34		12	8%
838.0	I	Closed dislocation	132	28	21%	94	71%	7	23	31	53	92	124	150	195	262	365	92	113.94	27		10	8%
<b>838.03</b>		<b>Closed dislocation</b>	<b>32</b>		<b>0%</b>	<b>31</b>	<b>97%</b>	<b>30</b>	<b>52</b>	<b>92</b>	<b>117</b>	<b>132</b>	<b>150</b>	<b>171</b>	<b>198</b>	<b>309</b>	<b>365</b>	<b>132</b>	<b>145.35</b>	<b>27</b>		<b>1</b>	<b>3%</b>
<b>838.09</b>		<b>Closed dislocation</b>	<b>40</b>	<b>22</b>	<b>55%</b>	<b>16</b>	<b>40%</b>	<b>1</b>	<b>5</b>	<b>5</b>	<b>18</b>	<b>23</b>	<b>25</b>	<b>31</b>	<b>47</b>	<b>89</b>	<b>201</b>	<b>23</b>	<b>35.69</b>	<b>27</b>		<b>2</b>	<b>5%</b>
839	I	Other, multiple, and ill-defined dislocations	171	69	40%	87	51%	1	2	4	5	8	17	24	52	137	365	8	41.18	13		15	9%
839.2	I	Thoracic and lumbar vertebra, closed	57	26	46%	28	49%	1	2	4	4	5	6	10	17	30	137	5	15.75	15		3	5%
<b>839.20</b>		<b>Lumbar vertebra</b>	<b>40</b>	<b>18</b>	<b>45%</b>	<b>20</b>	<b>50%</b>	<b>1</b>	<b>2</b>	<b>4</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>6</b>	<b>10</b>	<b>17</b>	<b>137</b>	<b>5</b>	<b>13.30</b>	<b>6</b>		<b>2</b>	<b>5%</b>
839.4	I	Other vertebra, closed	42	16	38%	21	50%	1	1	3	4	6	12	17	35	81	254	6	35.95	6		5	12%
839.6	I	Other location, closed	58	23	40%	29	50%	2	2	3	5	17	24	39	57	86	365	17	37.34	17		6	10%
<b>839.69</b>		<b>Other</b>	<b>43</b>	<b>19</b>	<b>44%</b>	<b>20</b>	<b>47%</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>9</b>	<b>17</b>	<b>24</b>	<b>39</b>	<b>89</b>	<b>4</b>	<b>17.05</b>	<b>5</b>		<b>4</b>	<b>9%</b>
840	I	Sprains and strains of shoulder and upper arm	39,714	21,585	54%	16,316	41%	1	2	3	4	6	11	23	49	125	365	6	39.61	8	17	1,813	5%
<b>840.0</b>		<b>Acromioclavicular (joint) (ligament)</b>	<b>1,018</b>	<b>631</b>	<b>62%</b>	<b>362</b>	<b>36%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>8</b>	<b>14</b>	<b>30</b>	<b>65</b>	<b>365</b>	<b>5</b>	<b>23.38</b>	<b>5</b>		<b>25</b>	<b>2%</b>
<b>840.3</b>		<b>Infraspinatus (muscle) (tendon)</b>	<b>87</b>	<b>26</b>	<b>30%</b>	<b>55</b>	<b>63%</b>	<b>3</b>	<b>6</b>	<b>19</b>	<b>37</b>	<b>56</b>	<b>80</b>	<b>127</b>	<b>200</b>	<b>365</b>	<b>365</b>	<b>56</b>	<b>107.11</b>	<b>15</b>		<b>6</b>	<b>7%</b>
<b>840.4</b>		<b>Rotator cuff (capsule)</b>	<b>3,772</b>	<b>1,264</b>	<b>34%</b>	<b>2,150</b>	<b>57%</b>	<b>3</b>	<b>8</b>	<b>25</b>	<b>51</b>	<b>84</b>	<b>112</b>	<b>153</b>	<b>210</b>	<b>324</b>	<b>365</b>	<b>84</b>	<b>115.60</b>	<b>16</b>	<b>20</b>	<b>358</b>	<b>9%</b>
<b>840.5</b>		<b>Subscapularis (muscle)</b>	<b>88</b>	<b>49</b>	<b>56%</b>	<b>35</b>	<b>40%</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>8</b>	<b>12</b>	<b>24</b>	<b>42</b>	<b>115</b>	<b>365</b>	<b>8</b>	<b>38.14</b>	<b>6</b>		<b>4</b>	<b>5%</b>
<b>840.6</b>		<b>Supraspinatus (muscle) (tendon)</b>	<b>387</b>	<b>137</b>	<b>35%</b>	<b>203</b>	<b>52%</b>	<b>2</b>	<b>2</b>	<b>5</b>	<b>14</b>	<b>34</b>	<b>97</b>	<b>129</b>	<b>200</b>	<b>277</b>	<b>365</b>	<b>34</b>	<b>97.24</b>	<b>5</b>		<b>47</b>	<b>12%</b>
<b>840.7</b>		<b>Superior glenoid labrum lesion</b>	<b>218</b>	<b>25</b>	<b>11%</b>	<b>150</b>	<b>69%</b>	<b>20</b>	<b>52</b>	<b>82</b>	<b>119</b>	<b>152</b>	<b>192</b>	<b>262</b>	<b>344</b>	<b>365</b>	<b>365</b>	<b>152</b>	<b>178.11</b>	<b>18</b>		<b>43</b>	<b>20%</b>
<b>840.8</b>		<b>Other specified sites of shoulder and upper arm</b>	<b>3,771</b>	<b>2,168</b>	<b>57%</b>	<b>1,490</b>	<b>40%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>17</b>	<b>48</b>	<b>365</b>	<b>3</b>	<b>19.78</b>	<b>4</b>		<b>113</b>	<b>3%</b>
<b>840.9</b>		<b>Unspecified site of shoulder and upper arm</b>	<b>30,346</b>	<b>17,268</b>	<b>57%</b>	<b>11,862</b>	<b>39%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>14</b>	<b>28</b>	<b>66</b>	<b>365</b>	<b>5</b>	<b>25.79</b>	<b>7</b>		<b>1,216</b>	<b>4%</b>
841	I	Sprains and strains of elbow and forearm	8,453	5,319	63%	2,835	34%	1	2	2	3	4	6	13	25	54	365	4	22.48	7		299	4%
<b>841.0</b>		<b>Radial collateral ligament</b>	<b>118</b>	<b>67</b>	<b>57%</b>	<b>41</b>	<b>35%</b>	<b>2</b>	<b>4</b>	<b>7</b>	<b>24</b>	<b>42</b>	<b>50</b>	<b>79</b>	<b>153</b>	<b>278</b>	<b>365</b>	<b>42</b>	<b>84.44</b>	<b>24</b>		<b>10</b>	<b>8%</b>
<b>841.1</b>		<b>Ulnar collateral ligament</b>	<b>124</b>	<b>49</b>	<b>40%</b>	<b>65</b>	<b>52%</b>	<b>2</b>	<b>3</b>	<b>6</b>	<b>11</b>	<b>18</b>	<b>41</b>	<b>51</b>	<b>82</b>	<b>179</b>	<b>365</b>	<b>18</b>	<b>63.55</b>	<b>18</b>		<b>10</b>	<b>8%</b>
<b>841.2</b>		<b>Radiohumeral (joint)</b>	<b>80</b>	<b>56</b>	<b>70%</b>	<b>20</b>	<b>25%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>8</b>	<b>44</b>	<b>2</b>	<b>5.25</b>	<b>3</b>		<b>4</b>	<b>5%</b>
<b>841.8</b>		<b>Other specified sites of elbow and forearm</b>	<b>414</b>	<b>257</b>	<b>62%</b>	<b>145</b>	<b>35%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>4</b>	<b>5</b>	<b>10</b>	<b>17</b>	<b>37</b>	<b>107</b>	<b>365</b>	<b>5</b>	<b>35.81</b>	<b>6</b>		<b>12</b>	<b>3%</b>
<b>841.9</b>		<b>Unspecified site of elbow and forearm</b>	<b>7,709</b>	<b>4,886</b>	<b>63%</b>	<b>2,561</b>	<b>33%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>12</b>	<b>22</b>	<b>47</b>	<b>365</b>	<b>4</b>	<b>19.83</b>	<b>5</b>		<b>262</b>	<b>3%</b>
842	I	Sprains and strains of wrist and hand	30,723	21,184	69%	8,655	28%	1	1	2	2	3	5	7	15	37	365	3	16.92	6	14	884	3%
842.0	I	Wrist	18,498	12,350	67%	5,572	30%	1	1	2	3	3	5	7	16	37	365	3	17.48	4		576	3%
<b>842.00</b>		<b>Unspecified site</b>	<b>17,403</b>	<b>11,780</b>	<b>68%</b>	<b>5,110</b>	<b>29%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>6</b>	<b>13</b>	<b>30</b>	<b>365</b>	<b>3</b>	<b>13.46</b>	<b>4</b>		<b>513</b>	<b>3%</b>
<b>842.01</b>		<b>Carpal (joint)</b>	<b>284</b>	<b>162</b>	<b>57%</b>	<b>108</b>	<b>38%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>13</b>	<b>21</b>	<b>34</b>	<b>77</b>	<b>365</b>	<b>7</b>	<b>29.99</b>	<b>6</b>		<b>14</b>	<b>5%</b>
<b>842.02</b>		<b>Radiocarpal (joint) (ligament)</b>	<b>325</b>	<b>221</b>	<b>68%</b>	<b>97</b>	<b>30%</b>	<b>1</b>	<b>2</b>	<b>2</b>													

### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open	
842.12		Metacarpophalangeal (joint)	527	317	60%	188	36%	1	2	3	5	13	19	32	63	132	365	13	42.66	10		22	4%	
842.13		Interphalangeal (joint)	904	682	75%	206	23%	1	1	2	3	3	5	7	18	44	365	3	17.18	4		16	2%	
842.19		Other	194	147	76%	38	20%	1	1	2	2	4	5	10	15	23	241	4	15.13	5		9	5%	
843	I	Sprains and strains of hip and thigh	5,659	2,933	52%	2,498	44%	1	2	2	4	5	8	14	27	62	365	5	24.26	8		228	4%	
843.0		Iliofemoral (ligament)	316	199	63%	103	33%	1	2	3	4	6	10	15	27	56	365	6	25.26	6		14	4%	
843.8		Other specified sites of hip and thigh	901	475	53%	393	44%	1	2	2	5	8	13	20	29	60	365	8	23.36	8		33	4%	
843.9		Unspecified site of hip and thigh	4,431	2,256	51%	1,995	45%	1	2	2	3	5	7	13	25	63	365	5	24.45	4		180	4%	
844	I	Sprains and strains of knee and leg	25,197	12,769	51%	11,472	46%	1	2	3	4	5	10	18	34	78	365	5	28.33	11	12	956	4%	
844.0		Lateral collateral ligament of knee	382	204	53%	166	43%	1	2	3	4	6	12	24	44	84	365	6	32.73	6		12	3%	
844.1		Medial collateral ligament of knee	1,189	561	47%	573	48%	1	2	4	6	12	21	36	54	86	365	12	34.08	15		55	5%	
844.2		Cruciate ligament of knee	1,245	219	18%	906	73%	7	25	42	63	86	114	150	201	295	365	86	117.49	15		120	10%	
844.3		Tibiofibular (joint) (ligament), superior	43	23	53%	15	35%	2	3	17	18	22	29	44	76	222	259	22	62.07	21		5	12%	
844.8		Other specified sites of knee and leg	1,374	630	46%	702	51%	1	2	3	5	9	17	28	48	104	365	9	37.29	7		42	3%	
844.9		Unspecified site of knee and leg	20,964	11,132	53%	9,110	43%	1	2	2	3	5	6	12	21	45	365	5	18.27	9		722	3%	
845	I	Sprains and strains of ankle and foot	33,815	17,593	52%	15,173	45%	1	2	2	3	4	5	7	14	29	365	4	13.02	7	7	1,049	3%	
845.0	I	Ankle	27,520	14,048	51%	12,613	46%	1	2	2	3	4	5	8	14	30	365	4	13.21	6		859	3%	
845.00		Unspecified site	25,862	13,318	51%	11,763	45%	1	2	2	3	4	5	7	13	26	365	4	11.57	4		781	3%	
845.01		Deltoid (ligament), ankle	198	97	49%	96	48%	1	2	3	4	5	11	19	30	62	152	5	19.20	11		5	3%	
845.02		Calcaneofibular (ligament)	318	159	50%	143	45%	2	2	3	4	6	7	12	23	47	341	6	20.64	7		16	5%	
845.03		Tibiofibular (ligament), distal	191	77	40%	107	56%	2	4	9	13	20	31	51	74	139	365	20	50.14	14		7	4%	
845.09		Other	951	397	42%	504	53%	1	2	4	6	11	19	37	64	118	365	11	40.57	10		50	5%	
845.1	I	Foot	6,295	3,545	56%	2,560	41%	1	1	2	3	3	4	6	12	26	365	3	12.05	4		190	3%	
845.10		Unspecified site	5,968	3,348	56%	2,440	41%	1	1	2	3	3	4	6	12	25	365	3	11.84	4		180	3%	
845.11		Tarsometatarsal (joint) (ligament)	102	65	64%	32	31%	1	1	2	3	3	5	6	12	47	85	3	12.28	4		5	5%	
845.12		Metatarsophalangeal (joint)	57	33	58%	24	42%	1	1	3	3	4	10	14	20	41	365	4	26.17	6			0%	
845.19		Other	109	55	50%	51	47%	1	2	2	2	3	4	5	9	58	160	3	16.12	3		3	3%	
846	I	Sprains and strains of sacroiliac region	37,008	16,755	45%	18,482	50%	1	2	3	4	5	7	13	24	55	365	5	24.12	8	8	1,771	5%	
846.0		Lumbosacral (joint) (ligament)	31,994	14,325	45%	16,100	50%	1	2	3	4	5	8	14	25	58	365	5	24.73	5		1,569	5%	
846.1		Sacroiliac ligament	2,140	1,053	49%	999	47%	1	2	3	4	6	10	15	25	54	365	6	25.49	6		88	4%	
846.2		Sacrospinal (ligament)	36	14	39%	19	53%	1	1	2	3	6	6	16	37	49	190	6	21.05	4		3	8%	
846.8		Other specified sites of sacroiliac region	75	35	47%	39	52%	1	1	2	3	4	5	9	12	24	99	4	11.64	4		1	1%	
846.9		Unspecified site of sacroiliac region	2,748	1,321	48%	1,317	48%	1	2	2	3	4	5	6	12	31	365	4	15.98	4		110	4%	
847	I	Sprains and strains of other and unspecified parts of back	105,693	51,047	48%	49,680	47%	1	2	2	3	5	6	11	21	52	365	5	22.80	10		7	4,966	5%
847.0		Neck	35,319	17,404	49%	16,179	46%	1	2	2	3	5	8	15	29	70	365	5	27.67	5	8	1,736	5%	
847.1		Thoracic	19,245	10,207	53%	8,116	42%	1	2	2	3	4	5	8	16	40	365	4	18.99	9	6	922	5%	
847.2		Lumbar sprains and strains	50,344	23,052	46%	25,020	50%	1	2	2	3	5	6	10	19	44	365	5	20.78	9	8	2,272	5%	
847.3		Sacrum	589	281	48%	286	49%	1	2	3	4	6	9	15	30	53	365	6	30.18	6	7	22	4%	
847.4		Coccyx	194	102	53%	78	40%	1	2	4	6	6	10	19	32	72	342	6	28.54	6	9	14	7%	
848	I	Other and ill-defined sprains and strains	9,990	5,856	59%	3,666	37%	1	2	2	3	4	5	7	13	26	365	4	12.25	6	10	468	5%	
848.1		Jaw	63	43	68%	18	29%	1	1	1	2	2	3	4	5	30	36	2	5.89	3		2	3%	
848.3		Ribs	1,456	883	61%	501	34%	1	1	2	3	3	4	5	8	17	248	3	7.28	4		72	5%	
848.4	I	Sternum	301	176	58%	107	36%	1	2	3	3	4	6	8	16	36	365	4	22.07	5		18	6%	
848.40		Unspecified site	127	71	56%	48	38%	1	2	3	3	4	5	7	13	17	282	4	15.85	4		8	6%	
848.41		Sternoclavicular (joint) (ligament)	74	48	65%	21	28%	2	2	3	11	18	19	34	38	87	365	18	47.67	13		5	7%	
848.42		Chondrosternal (joint)	96	56	58%	35	36%	1	2	3	3	4	5	6	6	10	365	4	14.91	5		5	5%	
848.5		Pelvis	4,054	2,201	54%	1,653	41%	1	2	2	3	4	6	10	17	32	365	4	15.67	5		200	5%	
848.8		Other specified sites of sprains and strains	4,108	2,549	62%	1,383	34%	1	1	2	2	3	4	6	10	20	365	3	9.27	4		176	4%	
850	I	Concussion	2,796	1,178	42%	1,475	53%	1	1	2	3	4	5	8	16	35	365	4	17.67	5	5	143	5%	
850.0	I	With no loss of consciousness	1,002	499	50%	462	46%	1	1	1	2	3	3	5	7	18	351	3	8.20	3		41	4%	
850.1	I	With brief loss of consciousness	571	198	35%	343	60%	1	2	3	3	5	7	12	21	46	365	5	23.64	5		30	5%	
850.5		With loss of consciousness of unspecified duration	330	127	38%	185	56%	1	2	3	4	6	7	12	19	57	365	6	23.03	5		18	5%	

### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
<b>850.9</b>		<b>Concussion, unspecified</b>	<b>878</b>	<b>351</b>	<b>40%</b>	<b>474</b>	<b>54%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>10</b>	<b>21</b>	<b>43</b>	<b>365</b>	<b>4</b>	<b>20.13</b>	<b>4</b>		<b>53</b>	<b>6%</b>
851	I	Cerebral laceration and contusion	42	10	24%	27	64%	1	3	5	7	11	17	30	36	144	190	11	31.89	17		5	12%
851.0	I	Cortex (cerebral) contusion without mention of open intracranial wound	22	3	14%	16	73%	2	3	5	7	11	12	25	30	41	190	11	25.00	16		3	14%
852	I	Subarachnoid, subdural, and extradural hemorrhage, following injury	72	8	11%	48	67%	9	21	32	53	75	97	145	221	364	365	75	113.92	43	30	16	22%
852.2	I	Subdural hemorrhage following injury without mention of open intracranial w	45	6	13%	28	62%	7	16	36	58	84	111	171	235	365	365	84	127.04	17		11	24%
854	I	Intracranial injury of other and unspecified nature	882	361	41%	426	48%	1	1	2	3	5	9	17	27	77	365	5	31.52	6	2	95	11%
854.0	I	Without mention of open intracranial wound	832	347	42%	411	49%	1	1	2	3	4	8	16	26	65	365	4	28.49	5		74	9%
<b>854.01</b>		<b>Without mention of open intracranial wound</b>	<b>677</b>	<b>315</b>	<b>47%</b>	<b>325</b>	<b>48%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>12</b>	<b>19</b>	<b>35</b>	<b>365</b>	<b>4</b>	<b>20.25</b>	<b>5</b>		<b>37</b>	<b>5%</b>
<b>854.02</b>		<b>Without mention of open intracranial wound</b>	<b>94</b>	<b>20</b>	<b>21%</b>	<b>65</b>	<b>69%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>11</b>	<b>14</b>	<b>24</b>	<b>34</b>	<b>112</b>	<b>365</b>	<b>11</b>	<b>35.55</b>	<b>5</b>		<b>9</b>	<b>10%</b>
854.1	I	With open intracranial wound	50	14	28%	15	30%	6	6	9	9	20	61	182	239	365	365	20	114.47	9		21	42%
860	I	Traumatic pneumothorax and hemothorax	62	7	11%	51	82%	12	20	29	33	42	52	73	90	167	365	42	69.25	52		4	6%
<b>860.0</b>		<b>Pneumothorax without mention of open wound into thorax</b>	<b>40</b>	<b>3</b>	<b>8%</b>	<b>34</b>	<b>85%</b>	<b>9</b>	<b>13</b>	<b>23</b>	<b>29</b>	<b>30</b>	<b>41</b>	<b>52</b>	<b>73</b>	<b>91</b>	<b>355</b>	<b>30</b>	<b>52.50</b>	<b>52</b>		<b>3</b>	<b>8%</b>
861	I	Injury to heart and lung	81	9	11%	55	68%	8	20	29	39	51	56	88	108	164	365	51	73.25	43		17	21%
861.2	I	Lung, without mention of open wound into thorax	68	6	9%	51	75%	8	25	29	39	51	56	87	102	134	365	51	69.14	53		11	16%
<b>861.21</b>		<b>Contusion</b>	<b>66</b>	<b>6</b>	<b>9%</b>	<b>50</b>	<b>76%</b>	<b>8</b>	<b>25</b>	<b>29</b>	<b>39</b>	<b>51</b>	<b>56</b>	<b>87</b>	<b>102</b>	<b>134</b>	<b>365</b>	<b>51</b>	<b>70.40</b>	<b>56</b>		<b>10</b>	<b>15%</b>
863	I	Injury to gastrointestinal tract	49	20	41%	28	57%	2	3	7	45	52	98	132	167	365	365	52	104.93	7		1	2%
864	I	Injury to liver	28	3	11%	18	64%	6	11	17	32	40	47	58	73	83	365	40	55.83	40		7	25%
864.0	I	Without mention of open wound into cavity	28	3	11%	18	64%	6	11	17	32	40	47	58	73	83	365	40	55.83	41		7	25%
865	I	Injury to spleen	49	4	8%	40	82%	12	26	32	38	43	60	68	80	115	204	43	58.88	40		5	10%
865.0	I	Without mention of open wound into cavity	48	4	8%	39	81%	12	26	32	38	43	62	69	85	117	204	43	58.85	43		5	10%
866	I	Injury to kidney	43	14	33%	24	56%	2	3	5	6	19	26	35	57	97	365	19	46.67	19		5	12%
866.0	I	Without mention of open wound into cavity	42	13	31%	24	57%	2	3	5	6	19	26	35	57	97	365	19	46.67	22		5	12%
<b>866.01</b>		<b>Without mention of open wound into cavity</b>	<b>33</b>	<b>13</b>	<b>39%</b>	<b>17</b>	<b>52%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>19</b>	<b>20</b>	<b>31</b>	<b>257</b>	<b>365</b>	<b>6</b>	<b>48.29</b>	<b>22</b>		<b>3</b>	<b>9%</b>
867	I	Injury to pelvic organs	24	6	25%	15	63%	6	14	30	46	87	88	121	146	365	365	87	120.53	6		3	13%
868	I	Injury to other intra-abdominal organs	46	28	61%	16	35%	1	3	4	9	9	27	48	55	91	365	9	46.19	10		2	4%
868.0	I	Without mention of open wound into cavity	45	28	62%	15	33%	1	2	4	6	9	10	40	48	68	365	9	43.20	10		2	4%
870	I	Open wound of ocular adnexa	806	608	75%	177	22%	1	1	2	2	3	4	5	6	13	364	3	8.36	6		21	3%
<b>870.0</b>		<b>Laceration of skin of eyelid and periocular area</b>	<b>263</b>	<b>200</b>	<b>76%</b>	<b>53</b>	<b>20%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>11</b>	<b>80</b>	<b>3</b>	<b>5.38</b>	<b>3</b>		<b>10</b>	<b>4%</b>
<b>870.8</b>		<b>Other specified open wounds of ocular adnexa</b>	<b>455</b>	<b>353</b>	<b>78%</b>	<b>92</b>	<b>20%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>364</b>	<b>3</b>	<b>8.87</b>	<b>4</b>		<b>10</b>	<b>2%</b>
871	I	Open wound of eyeball	736	491	67%	217	29%	1	1	2	3	4	4	6	13	31	267	4	13.86	4		28	4%
<b>871.0</b>		<b>Ocular laceration without prolapse of intraocular tissue</b>	<b>60</b>	<b>24</b>	<b>40%</b>	<b>33</b>	<b>55%</b>	<b>1</b>	<b>2</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>12</b>	<b>16</b>	<b>20</b>	<b>30</b>	<b>174</b>	<b>6</b>	<b>15.55</b>	<b>5</b>		<b>3</b>	<b>5%</b>
<b>871.4</b>		<b>Unspecified laceration of eye</b>	<b>340</b>	<b>233</b>	<b>69%</b>	<b>93</b>	<b>27%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>12</b>	<b>108</b>	<b>3</b>	<b>7.43</b>	<b>3</b>		<b>14</b>	<b>4%</b>
<b>871.6</b>		<b>Penetration of eyeball with (nonmagnetic) foreign body</b>	<b>190</b>	<b>150</b>	<b>79%</b>	<b>36</b>	<b>19%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>267</b>	<b>2</b>	<b>10.58</b>	<b>2</b>		<b>4</b>	<b>2%</b>
<b>871.9</b>		<b>Unspecified open wound of eyeball</b>	<b>67</b>	<b>43</b>	<b>64%</b>	<b>23</b>	<b>34%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>73</b>	<b>2</b>	<b>5.83</b>	<b>4</b>		<b>1</b>	<b>1%</b>
872	I	Open wound of ear	779	607	78%	156	20%	1	1	2	2	3	4	6	9	16	365	3	10.35	3		16	2%
872.0	I	External ear, without mention of complication	319	254	80%	59	18%	1	1	2	2	3	4	6	10	16	365	3	11.53	4		6	2%
<b>872.00</b>		<b>External ear, unspecified site</b>	<b>188</b>	<b>152</b>	<b>81%</b>	<b>36</b>	<b>19%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>4</b>	<b>10</b>	<b>13</b>	<b>365</b>	<b>3</b>	<b>15.06</b>	<b>3</b>			<b>0%</b>
<b>872.01</b>		<b>Auricle, ear</b>	<b>120</b>	<b>93</b>	<b>78%</b>	<b>21</b>	<b>18%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>9</b>	<b>16</b>	<b>20</b>	<b>5</b>	<b>6.43</b>	<b>5</b>		<b>6</b>	<b>5%</b>
872.6	I	Other specified parts of ear, without mention of complication	72	51	71%	20	28%	1	1	1	2	2	3	5	6	20	23	2	5.50	5		1	1%
<b>872.61</b>		<b>Ear drum</b>	<b>53</b>	<b>36</b>	<b>68%</b>	<b>17</b>	<b>32%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>5</b>	<b>6</b>	<b>6</b>	<b>20</b>	<b>23</b>	<b>2</b>	<b>6.00</b>	<b>6</b>			<b>0%</b>
<b>872.8</b>		<b>Ear, part unspecified, without mention of complication</b>	<b>352</b>	<b>280</b>	<b>80%</b>	<b>64</b>	<b>18%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>14</b>	<b>365</b>	<b>2</b>	<b>11.20</b>	<b>4</b>		<b>8</b>	<b>2%</b>
873	I	Other open wound of head	22,052	17,758	81%	3,727	17%	1	1	1	2	2	3	4	5	9	365	2	7.24	3		567	3%
<b>873.0</b>		<b>Scalp, without mention of complication</b>	<b>6,626</b>	<b>5,312</b>	<b>80%</b>	<b>1,165</b>	<b>18%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>365</b>	<b>2</b>	<b>5.56</b>	<b>2</b>		<b>149</b>	<b>2%</b>
<b>873.1</b>		<b>Scalp, complicated</b>	<b>63</b>	<b>41</b>	<b>65%</b>	<b>20</b>	<b>32%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>12</b>	<b>16</b>	<b>21</b>	<b>60</b>	<b>253</b>	<b>4</b>	<b>25.15</b>	<b>4</b>		<b>2</b>	<b>3%</b>
873.2	I	Nose, without mention of complication	1,049	853	81%	171	16%	1	1	1	2	2	3	3	5	7	365	2	8.41	3		25	2%
<b>873.20</b>		<b>Nose, unspecified site</b>	<b>1,016</b>	<b>827</b>	<b>81%</b>	<b>165</b>	<b>16%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>365</b>	<b>2</b>	<b>8.46</b>	<b>3</b>		<b>24</b>	<b>2%</b>
873.4	I	Face, without mention of complication	10,517	8,656	82%	1,594	15%	1	1	1	2	2	3	4	5	8	365	2	5.46	3		267	3%
<b>873.40</b>		<b>Face, unspecified site</b>	<b>1,627</b>	<b>1,346</b>	<b>83%</b>	<b>244</b>	<b>15%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>37</b>	<b>2</b>	<b>3.41</b>	<b>3</b>		<b>37</b>	<b>2%</b>
<b>873.41</b>		<b>Cheek</b>	<b>743</b>	<b>621</b>	<b>84%</b>	<b>108</b>	<b>15%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>9</b>	<b>170</b>	<b>2</b>	<b>5.52</b>	<b>3</b>		<b>14</b>	<b>2%</b>
<b>873.42</b>		<b>Forehead</b>	<b>5,195</b>	<b>4,281</b>	<b>82%</b>	<b>778</b>	<b>15%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>6</b>	<b>365</b>	<b>2</b>	<b>6.13</b>	<b>2</b>		<b>136</b>	<b>3%</b>
<b>873.43</b>		<b>Lip</b>	<b>1,743</b>	<b>1,427</b>	<b>82%</b>	<b>268</b>	<b>15%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>											

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
873.44		<b>Jaw</b>	<b>1,001</b>	<b>846</b>	<b>85%</b>	<b>129</b>	<b>13%</b>	1	1	1	2	2	3	4	6	10	80	2	4.87	3		26	3%
873.49		<b>Other and multiple sites</b>	<b>208</b>	<b>135</b>	<b>65%</b>	<b>67</b>	<b>32%</b>	1	2	2	2	3	5	8	11	18	34	3	6.90	4		6	3%
873.5	I	Face, complicated	258	167	65%	82	32%	1	2	2	3	4	5	7	13	32	157	4	13.40	4		9	3%
873.52		<b>Forehead</b>	<b>70</b>	<b>45</b>	<b>64%</b>	<b>22</b>	<b>31%</b>	1	1	2	3	4	13	24	32	41	75	4	16.86	4		3	4%
873.53		<b>Lip</b>	<b>60</b>	<b>33</b>	<b>55%</b>	<b>24</b>	<b>40%</b>	1	1	2	3	3	4	4	5	7	13	3	3.83	4		3	5%
873.6	I	Internal structures of mouth, without mention of complication	2,254	1,720	76%	455	20%	1	1	2	2	3	3	5	6	16	365	3	13.52	3		79	4%
873.60		<b>Mouth, unspecified site</b>	<b>89</b>	<b>70</b>	<b>79%</b>	<b>15</b>	<b>17%</b>	1	2	3	4	5	6	6	8	10	86	5	10.20	4		4	4%
873.63		<b>Tooth (broken)</b>	<b>2,051</b>	<b>1,574</b>	<b>77%</b>	<b>405</b>	<b>20%</b>	1	1	2	2	2	3	4	6	15	365	2	12.20	3		72	4%
873.7	I	Internal structures of mouth, complicated	111	78	70%	30	27%	1	2	3	4	10	14	20	25	139	314	10	36.50	5		3	3%
873.73		<b>Tooth (broken)</b>	<b>94</b>	<b>69</b>	<b>73%</b>	<b>24</b>	<b>26%</b>	1	1	2	3	4	14	17	25	139	314	4	32.96	4		1	1%
873.8		<b>Other and unspecified open wound of head without mention of complic</b>	<b>1,134</b>	<b>900</b>	<b>79%</b>	<b>202</b>	<b>18%</b>	1	1	1	2	2	2	3	4	8	365	2	7.28	2		32	3%
874	I	Open wound of neck	234	174	74%	55	24%	1	1	2	2	3	4	5	8	24	92	3	9.47	4		5	2%
874.8		<b>Other and unspecified parts, without mention of complication</b>	<b>210</b>	<b>159</b>	<b>76%</b>	<b>46</b>	<b>22%</b>	1	2	2	2	2	4	5	8	24	92	2	8.72	4		5	2%
875	I	Open wound of chest (wall)	334	234	70%	84	25%	1	1	2	3	4	6	7	18	32	178	4	13.68	7		16	5%
875.0		<b>Without mention of complication</b>	<b>291</b>	<b>209</b>	<b>72%</b>	<b>70</b>	<b>24%</b>	1	1	2	2	3	5	7	11	30	178	3	11.77	5		12	4%
876	I	Open wound of back	151	120	79%	29	19%	1	1	1	2	2	2	3	5	6	25	2	3.28	3		2	1%
876.0		<b>Without mention of complication</b>	<b>142</b>	<b>114</b>	<b>80%</b>	<b>26</b>	<b>18%</b>	1	1	1	2	2	2	3	4	6	25	2	3.31	2		2	1%
877	I	Open wound of buttock	179	131	73%	42	23%	1	2	2	3	4	6	8	18	25	71	4	10.40	4		6	3%
877.0		<b>Without mention of complication</b>	<b>170</b>	<b>127</b>	<b>75%</b>	<b>37</b>	<b>22%</b>	1	2	2	3	4	6	7	18	25	42	4	9.05	4		6	4%
878	I	Open wound of genital organs (external), including traumatic amputation	66	32	48%	32	48%	1	1	3	5	10	13	14	25	32	87	10	15.63	9		2	3%
878.2		<b>Scrotum and testes, without mention of complication</b>	<b>41</b>	<b>20</b>	<b>49%</b>	<b>20</b>	<b>49%</b>	1	1	1	2	5	7	10	13	25	87	5	12.80	5		1	2%
879	I	Open wound of other and unspecified sites, except limbs	506	379	75%	105	21%	1	1	2	3	3	5	10	17	32	95	3	11.20	8	5	22	4%
879.2		<b>Abdominal wall, anterior, without mention of complication</b>	<b>296</b>	<b>220</b>	<b>74%</b>	<b>61</b>	<b>21%</b>	1	1	2	2	3	3	4	6	17	82	3	6.90	3		15	5%
879.4		<b>Abdominal wall, lateral, without mention of complication</b>	<b>66</b>	<b>42</b>	<b>64%</b>	<b>20</b>	<b>30%</b>	1	2	4	5	7	10	16	19	33	95	7	17.20	10		4	6%
880	I	Open wound of shoulder and upper arm	1,348	1,103	82%	211	16%	1	1	2	2	3	4	6	12	28	365	3	14.00	4		34	3%
880.0	I	Without mention of complication	1,273	1,053	83%	188	15%	1	1	2	2	3	4	5	10	25	365	3	12.14	3		32	3%
880.00		<b>Without mention of complication</b>	<b>355</b>	<b>276</b>	<b>78%</b>	<b>61</b>	<b>17%</b>	1	1	2	3	4	9	10	20	33	111	4	14.13	3		18	5%
880.03		<b>Without mention of complication</b>	<b>877</b>	<b>748</b>	<b>85%</b>	<b>115</b>	<b>13%</b>	1	1	2	2	2	3	4	5	11	365	2	9.08	3		14	2%
880.1	I	Complicated	65	47	72%	17	26%	2	3	3	4	5	21	26	26	54	148	5	21.59	13		1	2%
881	I	Open wound of elbow, forearm, and wrist	18,196	15,026	83%	2,734	15%	1	1	2	2	3	3	5	8	19	365	3	9.87	4		436	2%
881.0	I	Without mention of complication	17,348	14,552	84%	2,399	14%	1	1	2	2	2	3	4	6	12	365	2	6.99	3		397	2%
881.00		<b>Without mention of complication</b>	<b>10,729</b>	<b>9,087</b>	<b>85%</b>	<b>1,370</b>	<b>13%</b>	1	1	2	2	2	3	4	6	12	365	2	6.69	3		272	3%
881.01		<b>Without mention of complication</b>	<b>2,272</b>	<b>1,857</b>	<b>82%</b>	<b>377</b>	<b>17%</b>	1	1	2	2	2	3	4	6	12	364	2	7.02	3		38	2%
881.02		<b>Without mention of complication</b>	<b>4,347</b>	<b>3,608</b>	<b>83%</b>	<b>652</b>	<b>15%</b>	1	1	1	2	2	3	4	6	11	364	2	7.61	3		87	2%
881.1	I	Complicated	573	385	67%	163	28%	1	2	3	4	6	9	14	25	40	200	6	14.06	9		25	4%
881.10		<b>Complicated</b>	<b>351</b>	<b>246</b>	<b>70%</b>	<b>88</b>	<b>25%</b>	1	2	3	4	6	7	14	23	41	200	6	14.03	9		17	5%
881.11		<b>Complicated</b>	<b>75</b>	<b>43</b>	<b>57%</b>	<b>28</b>	<b>37%</b>	1	3	6	9	9	13	32	34	46	69	9	19.11	9		4	5%
881.12		<b>Complicated</b>	<b>147</b>	<b>96</b>	<b>65%</b>	<b>47</b>	<b>32%</b>	1	1	2	3	5	7	9	15	28	76	5	11.11	9		4	3%
881.2	I	With tendon involvement	275	89	32%	172	63%	2	4	8	14	22	33	48	71	111	365	22	46.00	19		14	5%
881.20		<b>With tendon involvement</b>	<b>111</b>	<b>30</b>	<b>27%</b>	<b>76</b>	<b>68%</b>	5	10	16	20	30	43	60	75	120	365	30	52.71	19		5	5%
881.22		<b>With tendon involvement</b>	<b>157</b>	<b>57</b>	<b>36%</b>	<b>92</b>	<b>59%</b>	1	3	5	8	17	26	41	49	89	288	17	36.75	19		8	5%
882	I	Open wound of hand except finger(s) alone	29,418	23,855	81%	4,959	17%	1	1	2	2	3	3	5	8	17	365	3	8.97	4		604	2%
882.0		<b>Without mention of complication</b>	<b>27,856</b>	<b>22,972</b>	<b>82%</b>	<b>4,331</b>	<b>16%</b>	1	1	1	2	2	3	4	6	12	365	2	6.84	3		553	2%
882.1		<b>Complicated</b>	<b>1,088</b>	<b>703</b>	<b>65%</b>	<b>355</b>	<b>33%</b>	1	2	2	3	4	6	10	16	32	365	4	14.28	6		30	3%
882.2		<b>With tendon involvement</b>	<b>474</b>	<b>180</b>	<b>38%</b>	<b>273</b>	<b>58%</b>	2	3	6	12	18	25	39	58	90	365	18	35.77	20		21	4%
883	I	Open wound of finger(s)	104,948	85,491	81%	17,078	16%	1	1	2	2	2	3	5	9	21	365	2	10.16	5		2,379	2%
883.0		<b>Without mention of complication</b>	<b>98,575</b>	<b>82,054</b>	<b>83%</b>	<b>14,342</b>	<b>15%</b>	1	1	1	2	2	3	4	6	12	365	2	6.89	3		2,179	2%
883.1		<b>Complicated</b>	<b>3,522</b>	<b>2,337</b>	<b>66%</b>	<b>1,098</b>	<b>31%</b>	1	2	3	4	6	10	16	26	45	365	6	17.44	10		87	2%
883.2		<b>With tendon involvement</b>	<b>2,851</b>	<b>1,100</b>	<b>39%</b>	<b>1,638</b>	<b>57%</b>	2	4	6	12	17	26	40	55	83	365	17	33.86	16		113	4%
885	I	Traumatic amputation of thumb (complete) (partial)	248	44	18%	185	75%	4	6	11	17	22	37	47	68	103	365	22	49.10	27		19	8%
885.0		<b>Without mention of complication</b>	<b>239</b>	<b>42</b>	<b>18%</b>	<b>179</b>	<b>75%</b>	4	6	11	18	24	37	47	76	103	365	24	49.27	27		18	8%

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open	
886	I	Traumatic amputation of other finger(s) (complete) (partial)	1,984	338	17%	1,544	78%	3	6	12	18	27	38	48	66	108	365	27	46.47	28	17	102	5%	
<b>886.0</b>		<b>Without mention of complication</b>	<b>1,863</b>	<b>329</b>	<b>18%</b>	<b>1,436</b>	<b>77%</b>	<b>3</b>	<b>6</b>	<b>12</b>	<b>18</b>	<b>26</b>	<b>37</b>	<b>48</b>	<b>64</b>	<b>104</b>	<b>365</b>	<b>26</b>	<b>45.42</b>	<b>28</b>		<b>98</b>	<b>5%</b>	
<b>886.1</b>		<b>Complicated</b>	<b>121</b>	<b>9</b>	<b>7%</b>	<b>108</b>	<b>89%</b>	<b>6</b>	<b>11</b>	<b>15</b>	<b>24</b>	<b>33</b>	<b>45</b>	<b>63</b>	<b>83</b>	<b>162</b>	<b>365</b>	<b>33</b>	<b>60.51</b>	<b>35</b>		<b>4</b>	<b>3%</b>	
887	I	Traumatic amputation of arm and hand (complete) (partial)	36		0%	16	44%	102	153	184	196	244	365	365	365	365	365	244	257.25	153	54	20	56%	
890	I	Open wound of hip and thigh	2,445	1,930	79%	432	18%	1	1	2	2	3	4	5	10	20	365	3	10.30			83	3%	
<b>890.0</b>		<b>Without mention of complication</b>	<b>2,285</b>	<b>1,835</b>	<b>80%</b>	<b>371</b>	<b>16%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>7</b>	<b>15</b>	<b>365</b>	<b>3</b>	<b>9.07</b>	<b>3</b>		<b>79</b>	<b>3%</b>	
<b>890.1</b>		<b>Complicated</b>	<b>156</b>	<b>94</b>	<b>60%</b>	<b>58</b>	<b>37%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>9</b>	<b>14</b>	<b>23</b>	<b>41</b>	<b>222</b>	<b>5</b>	<b>17.22</b>	<b>7</b>		<b>4</b>	<b>3%</b>	
891	I	Open wound of knee, leg [except thigh], and ankle	8,125	6,157	76%	1,782	22%	1	1	2	2	3	4	5	10	20	365	3	9.85			186	2%	
<b>891.0</b>		<b>Without mention of complication</b>	<b>7,650</b>	<b>5,921</b>	<b>77%</b>	<b>1,566</b>	<b>20%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>14</b>	<b>365</b>	<b>2</b>	<b>7.04</b>	<b>3</b>		<b>163</b>	<b>2%</b>	
<b>891.1</b>		<b>Complicated</b>	<b>432</b>	<b>223</b>	<b>52%</b>	<b>191</b>	<b>44%</b>	<b>1</b>	<b>2</b>	<b>4</b>	<b>6</b>	<b>12</b>	<b>18</b>	<b>26</b>	<b>37</b>	<b>62</b>	<b>304</b>	<b>12</b>	<b>26.31</b>	<b>14</b>		<b>18</b>	<b>4%</b>	
<b>891.2</b>		<b>With tendon involvement</b>	<b>43</b>	<b>13</b>	<b>30%</b>	<b>25</b>	<b>58%</b>	<b>2</b>	<b>9</b>	<b>18</b>	<b>27</b>	<b>53</b>	<b>58</b>	<b>62</b>	<b>70</b>	<b>110</b>	<b>365</b>	<b>53</b>	<b>60.60</b>	<b>53</b>		<b>5</b>	<b>12%</b>	
892	I	Open wound of foot except toe(s) alone	4,043	3,042	75%	927	23%	1	1	2	2	3	3	4	6	12	365	3	7.40			74	2%	
<b>892.0</b>		<b>Without mention of complication</b>	<b>3,817</b>	<b>2,918</b>	<b>76%</b>	<b>829</b>	<b>22%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>11</b>	<b>365</b>	<b>3</b>	<b>6.77</b>	<b>3</b>		<b>70</b>	<b>2%</b>	
<b>892.1</b>		<b>Complicated</b>	<b>196</b>	<b>112</b>	<b>57%</b>	<b>81</b>	<b>41%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>6</b>	<b>8</b>	<b>17</b>	<b>91</b>	<b>3</b>	<b>7.69</b>	<b>5</b>		<b>3</b>	<b>2%</b>	
<b>892.2</b>		<b>With tendon involvement</b>	<b>30</b>	<b>12</b>	<b>40%</b>	<b>17</b>	<b>57%</b>	<b>1</b>	<b>1</b>	<b>5</b>	<b>10</b>	<b>14</b>	<b>28</b>	<b>30</b>	<b>43</b>	<b>121</b>	<b>225</b>	<b>14</b>	<b>36.41</b>	<b>21</b>		<b>1</b>	<b>3%</b>	
893	I	Open wound of toe(s)	865	573	66%	272	31%	1	1	2	3	4	5	6	11	21	315	4	10.11			20	2%	
<b>893.0</b>		<b>Without mention of complication</b>	<b>793</b>	<b>548</b>	<b>69%</b>	<b>229</b>	<b>29%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>9</b>	<b>17</b>	<b>315</b>	<b>3</b>	<b>9.03</b>	<b>5</b>		<b>16</b>	<b>2%</b>	
<b>893.1</b>		<b>Complicated</b>	<b>57</b>	<b>23</b>	<b>40%</b>	<b>30</b>	<b>53%</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>5</b>	<b>6</b>	<b>12</b>	<b>18</b>	<b>40</b>	<b>84</b>	<b>5</b>	<b>13.53</b>	<b>12</b>		<b>4</b>	<b>7%</b>	
895	I	Traumatic amputation of toe(s) (complete) (partial)	93	1	1%	69	74%	21	35	58	71	94	117	164	232	365	365	94	134.33			64	23	25%
<b>895.0</b>		<b>Without mention of complication</b>	<b>91</b>	<b>1</b>	<b>1%</b>	<b>67</b>	<b>74%</b>	<b>21</b>	<b>35</b>	<b>60</b>	<b>80</b>	<b>95</b>	<b>125</b>	<b>164</b>	<b>232</b>	<b>365</b>	<b>365</b>	<b>95</b>	<b>136.58</b>	<b>64</b>		<b>23</b>	<b>25%</b>	
903	I	Injury to blood vessels of upper extremity	142	14	10%	113	80%	8	14	22	31	52	69	87	123	329	365	52	93.30			35	15	11%
<b>903.2</b>		<b>Radial blood vessels</b>	<b>24</b>	<b>1</b>	<b>4%</b>	<b>22</b>	<b>92%</b>	<b>6</b>	<b>12</b>	<b>13</b>	<b>18</b>	<b>25</b>	<b>40</b>	<b>51</b>	<b>87</b>	<b>261</b>	<b>365</b>	<b>25</b>	<b>69.09</b>	<b>31</b>		<b>1</b>	<b>4%</b>	
<b>903.3</b>		<b>Ulnar blood vessels</b>	<b>30</b>	<b>3</b>	<b>10%</b>	<b>22</b>	<b>73%</b>	<b>17</b>	<b>32</b>	<b>59</b>	<b>69</b>	<b>98</b>	<b>172</b>	<b>281</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>98</b>	<b>164.91</b>	<b>33</b>		<b>5</b>	<b>17%</b>	
<b>903.5</b>		<b>Digital blood vessels</b>	<b>79</b>	<b>9</b>	<b>11%</b>	<b>62</b>	<b>78%</b>	<b>8</b>	<b>14</b>	<b>22</b>	<b>27</b>	<b>46</b>	<b>67</b>	<b>79</b>	<b>89</b>	<b>139</b>	<b>365</b>	<b>46</b>	<b>73.55</b>	<b>38</b>		<b>8</b>	<b>10%</b>	
910	I	Superficial injury of face, neck, and scalp except eye	3,570	2,714	76%	761	21%	1	1	1	2	2	3	4	5	8	365	2	5.60			95	3%	
<b>910.0</b>		<b>Abrasion or friction burn without mention of infection</b>	<b>3,156</b>	<b>2,370</b>	<b>75%</b>	<b>698</b>	<b>22%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>7</b>	<b>365</b>	<b>2</b>	<b>5.65</b>	<b>3</b>		<b>88</b>	<b>3%</b>	
<b>910.4</b>		<b>Insect bite, nonvenomous, without mention of infection</b>	<b>183</b>	<b>158</b>	<b>86%</b>	<b>23</b>	<b>13%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>2</b>	<b>1.87</b>	<b>3</b>			<b>2</b>	<b>1%</b>	
<b>910.6</b>		<b>Superficial foreign body (splinter) without major open wound and with</b>	<b>156</b>	<b>130</b>	<b>83%</b>	<b>22</b>	<b>14%</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>10</b>	<b>18</b>	<b>56</b>	<b>3</b>	<b>7.59</b>	<b>3</b>		<b>4</b>	<b>3%</b>	
911	I	Superficial injury of trunk	1,491	1,010	68%	438	29%	1	1	2	2	3	4	5	8	18	240	3	8.72			43	3%	
<b>911.0</b>		<b>Abrasion or friction burn without mention of infection</b>	<b>1,320</b>	<b>873</b>	<b>66%</b>	<b>410</b>	<b>31%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>8</b>	<b>19</b>	<b>240</b>	<b>3</b>	<b>9.10</b>	<b>3</b>		<b>37</b>	<b>3%</b>	
912	I	Superficial injury of shoulder and upper arm	901	641	71%	231	26%	1	1	2	2	2	3	5	7	12	195	2	7.51			29	3%	
<b>912.0</b>		<b>Abrasion or friction burn without mention of infection</b>	<b>767</b>	<b>540</b>	<b>70%</b>	<b>205</b>	<b>27%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>12</b>	<b>195</b>	<b>2</b>	<b>7.19</b>	<b>3</b>		<b>22</b>	<b>3%</b>	
913	I	Superficial injury of elbow, forearm, and wrist	4,654	3,622	78%	907	19%	1	1	2	2	2	3	4	19	16	364	2	7.97			125	3%	
<b>913.0</b>		<b>Abrasion or friction burn without mention of infection</b>	<b>3,978</b>	<b>3,094</b>	<b>78%</b>	<b>785</b>	<b>20%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>15</b>	<b>175</b>	<b>2</b>	<b>7.57</b>	<b>3</b>		<b>99</b>	<b>2%</b>	
<b>913.4</b>		<b>Insect bite, nonvenomous, without mention of infection</b>	<b>267</b>	<b>211</b>	<b>79%</b>	<b>46</b>	<b>17%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>9</b>	<b>364</b>	<b>2</b>	<b>12.85</b>	<b>3</b>		<b>10</b>	<b>4%</b>	
<b>913.6</b>		<b>Superficial foreign body (splinter) without major open wound and with</b>	<b>333</b>	<b>256</b>	<b>77%</b>	<b>66</b>	<b>20%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>11</b>	<b>15</b>	<b>25</b>	<b>78</b>	<b>4</b>	<b>9.74</b>	<b>3</b>		<b>11</b>	<b>3%</b>	
914	I	Superficial injury of hand(s) except finger(s) alone	3,254	2,556	79%	616	19%	1	1	2	2	2	3	4	6	12	116	2	5.49			82	3%	
<b>914.0</b>		<b>Abrasion or friction burn without mention of infection</b>	<b>2,295</b>	<b>1,819</b>	<b>79%</b>	<b>421</b>	<b>18%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>9</b>	<b>116</b>	<b>2</b>	<b>4.66</b>	<b>3</b>	<b>2</b>	<b>55</b>	<b>2%</b>	
<b>914.4</b>		<b>Insect bite, nonvenomous, without mention of infection</b>	<b>147</b>	<b>124</b>	<b>84%</b>	<b>19</b>	<b>13%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>2</b>	<b>2.11</b>	<b>3</b>		<b>4</b>	<b>3%</b>	
<b>914.6</b>		<b>Superficial foreign body (splinter) without major open wound and with</b>	<b>687</b>	<b>524</b>	<b>76%</b>	<b>146</b>	<b>21%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>10</b>	<b>17</b>	<b>84</b>	<b>3</b>	<b>7.31</b>	<b>4</b>	<b>1</b>	<b>17</b>	<b>2%</b>	
<b>914.7</b>		<b>Superficial foreign body (splinter) without major open wound, infected</b>	<b>45</b>	<b>28</b>	<b>62%</b>	<b>15</b>	<b>33%</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>8</b>	<b>13</b>	<b>17</b>	<b>24</b>	<b>38</b>	<b>78</b>	<b>8</b>	<b>16.27</b>	<b>6</b>		<b>2</b>	<b>4%</b>	
915	I	Superficial injury of finger(s)	4,922	4,063	83%	742	15%	1	1	2	2	2	3	4	5	11	317	2	6.29			117	2%	
<b>915.0</b>		<b>Abrasion or friction burn without mention of infection</b>	<b>2,715</b>	<b>2,254</b>	<b>83%</b>	<b>400</b>	<b>15%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>9</b>	<b>91</b>	<b>2</b>	<b>4.89</b>	<b>3</b>		<b>61</b>	<b>2%</b>	
<b>915.6</b>		<b>Superficial foreign body (splinter) without major open wound and with</b>	<b>1,954</b>	<b>1,610</b>	<b>82%</b>	<b>296</b>	<b>15%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>13</b>	<b>317</b>	<b>2</b>	<b>7.44</b>	<b>3</b>		<b>48</b>	<b>2%</b>	
<b>915.7</b>		<b>Superficial foreign body (splinter) without major open wound, infected</b>	<b>92</b>	<b>68</b>	<b>74%</b>	<b>22</b>	<b>24%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>6</b>	<b>10</b>	<b>19</b>	<b>24</b>	<b>3</b>	<b>6.27</b>	<b>4</b>		<b>2</b>	<b>2%</b>	
916	I	Superficial injury of hip, thigh, leg, and ankle	5,874	4,020	68%	1,672	28%	1	1	2	2	3	4	5	8	18	365	3	9.18			182	3%	
<b>916.0</b>		<b>Abrasion or friction burn without mention of infection</b>	<b>5,291</b>	<b>3,620&lt;/</b>																				

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
917.0		<b>Abrasion or friction burn without mention of infection</b>	<b>595</b>	<b>396</b>	<b>67%</b>	<b>184</b>	<b>31%</b>	1	1	2	2	3	4	6	9	19	95	3	8.52	4		15	3%
917.6		<b>Superficial foreign body (splinter) without major open wound and with</b>	<b>126</b>	<b>85</b>	<b>67%</b>	<b>37</b>	<b>29%</b>	1	2	3	3	4	5	5	7	18	93	4	8.30	4		4	3%
918	I	Superficial injury of eye and adnexa	22,329	16,740	75%	5,073	23%	1	1	1	1	2	2	3	4	5	365	2	3.87	2		516	2%
918.0		<b>Eyelids and periocular area</b>	<b>224</b>	<b>178</b>	<b>79%</b>	<b>35</b>	<b>16%</b>	1	1	1	2	2	2	2	3	4	16	2	2.49	2		11	5%
918.1		<b>Cornea</b>	<b>20,017</b>	<b>14,932</b>	<b>75%</b>	<b>4,634</b>	<b>23%</b>	1	1	1	1	2	2	3	3	5	365	2	3.75	2		451	2%
918.2		<b>Conjunctiva</b>	<b>1,086</b>	<b>856</b>	<b>79%</b>	<b>203</b>	<b>19%</b>	1	1	1	1	2	2	3	4	5	365	2	4.33	2		27	2%
918.9		<b>Other and unspecified superficial injuries of eye</b>	<b>1,002</b>	<b>774</b>	<b>77%</b>	<b>201</b>	<b>20%</b>	1	1	1	2	2	2	3	4	6	365	2	6.41	2		27	3%
920		<b>Contusion of face, scalp, and neck except eye(s)</b>	<b>13,304</b>	<b>9,720</b>	<b>73%</b>	<b>3,296</b>	<b>25%</b>	1	1	1	2	2	2	3	4	6	365	2	5.19	3		288	2%
921	I	Contusion of eye and adnexa	2,334	1,637	70%	647	28%	1	1	1	2	2	3	4	5	9	365	2	4.66	3		50	2%
921.0		<b>Black eye, NOS</b>	<b>104</b>	<b>69</b>	<b>66%</b>	<b>33</b>	<b>32%</b>	1	1	1	2	3	3	6	9	21	78	3	7.97	3		2	2%
921.1		<b>Contusion of eyelids and periocular area</b>	<b>367</b>	<b>264</b>	<b>72%</b>	<b>96</b>	<b>26%</b>	1	1	1	2	2	3	3	5	8	24	2	3.54	3		7	2%
921.2		<b>Contusion of orbital tissues</b>	<b>438</b>	<b>296</b>	<b>68%</b>	<b>134</b>	<b>31%</b>	1	1	1	2	2	3	3	4	6	77	2	3.85	2		8	2%
921.3		<b>Contusion of eyeball</b>	<b>414</b>	<b>270</b>	<b>65%</b>	<b>134</b>	<b>32%</b>	1	1	1	2	3	4	5	7	12	39	3	4.73	3		10	2%
921.9		<b>Unspecified contusion of eye</b>	<b>1,011</b>	<b>738</b>	<b>73%</b>	<b>250</b>	<b>25%</b>	1	1	1	2	2	3	4	6	6	365	2	5.05	2		23	2%
922	I	Contusion of trunk	19,954	11,662	58%	7,550	38%	1	1	2	2	3	4	5	9	19	365	3	10.00	5	7	742	4%
922.0		<b>Breast</b>	<b>223</b>	<b>154</b>	<b>69%</b>	<b>67</b>	<b>30%</b>	1	1	2	2	3	4	6	12	23	144	3	11.13	3		2	1%
922.1		<b>Chest wall</b>	<b>8,525</b>	<b>5,059</b>	<b>59%</b>	<b>3,105</b>	<b>36%</b>	1	1	2	2	3	4	6	9	19	365	3	9.07	4		361	4%
922.2		<b>Abdominal wall</b>	<b>1,994</b>	<b>1,101</b>	<b>55%</b>	<b>798</b>	<b>40%</b>	1	1	2	2	3	4	6	9	19	365	3	11.25	4		95	5%
922.3	I	Back	8,822	5,142	58%	3,407	39%	1	1	2	2	3	4	5	8	21	365	3	10.63	3		273	3%
922.31		<b>Back</b>	<b>6,810</b>	<b>3,928</b>	<b>58%</b>	<b>2,665</b>	<b>39%</b>	1	1	2	2	3	4	5	9	22	365	3	11.29	3		217	3%
922.32		<b>Buttock</b>	<b>2,002</b>	<b>1,208</b>	<b>60%</b>	<b>739</b>	<b>37%</b>	1	1	2	2	3	4	5	7	18	365	3	8.31	3		55	3%
922.4		<b>Genital organs</b>	<b>363</b>	<b>190</b>	<b>52%</b>	<b>162</b>	<b>45%</b>	1	1	2	2	3	4	5	7	11	40	3	5.09	3		11	3%
923	I	Contusion of upper limb	60,095	44,786	75%	13,820	23%	1	1	2	2	2	3	4	6	13	365	2	7.64	4		1,489	2%
923.0	I	Shoulder and upper arm	6,974	4,416	63%	2,372	34%	1	1	2	2	3	4	5	8	17	365	3	9.90	4	3	186	3%
923.00		<b>Shoulder region</b>	<b>5,352</b>	<b>3,349</b>	<b>63%</b>	<b>1,854</b>	<b>35%</b>	1	1	2	2	3	4	5	8	17	365	3	9.85	4		149	3%
923.01		<b>Scapular region</b>	<b>252</b>	<b>154</b>	<b>61%</b>	<b>89</b>	<b>35%</b>	1	1	2	2	3	4	5	11	27	365	3	13.02	4		9	4%
923.03		<b>Upper arm</b>	<b>1,354</b>	<b>903</b>	<b>67%</b>	<b>423</b>	<b>31%</b>	1	1	2	2	3	3	5	7	15	365	3	9.56	3		28	2%
923.1	I	Elbow and forearm	12,237	8,724	71%	3,196	26%	1	1	2	2	3	3	4	6	14	365	3	7.58	3		317	3%
923.10		<b>Forearm</b>	<b>3,861</b>	<b>2,895</b>	<b>75%</b>	<b>873</b>	<b>23%</b>	1	1	1	2	2	3	4	5	10	364	2	6.52	3		93	2%
923.11		<b>Elbow</b>	<b>8,376</b>	<b>5,829</b>	<b>70%</b>	<b>2,323</b>	<b>28%</b>	1	1	2	2	3	3	5	7	15	365	3	7.97	3		224	3%
923.2	I	Wrist and hand(s), except finger(s) alone	22,857	16,921	74%	5,363	23%	1	1	2	2	3	3	4	6	15	365	3	7.74	3	4	573	3%
923.20		<b>Hand(s)</b>	<b>15,396</b>	<b>11,710</b>	<b>76%</b>	<b>3,322</b>	<b>22%</b>	1	1	2	2	2	3	4	6	12	365	2	6.58	3		364	2%
923.21		<b>Wrist</b>	<b>7,461</b>	<b>5,211</b>	<b>70%</b>	<b>2,041</b>	<b>27%</b>	1	1	2	2	3	4	5	9	20	365	3	9.63	3		209	3%
923.3		<b>Finger</b>	<b>17,974</b>	<b>14,699</b>	<b>82%</b>	<b>2,865</b>	<b>16%</b>	1	1	1	2	2	3	3	5	8	365	2	5.62	2		410	2%
923.8		<b>Multiple sites of upper limb</b>	<b>52</b>	<b>25</b>	<b>48%</b>	<b>24</b>	<b>46%</b>	1	2	2	4	6	11	12	18	27	89	6	12.79	10		3	6%
924	I	Contusion of lower limb and of other and unspecified sites	47,598	29,990	63%	16,264	34%	1	1	2	2	3	4	5	8	18	365	3	9.12	6	9	1,344	3%
924.0	I	Hip and thigh	7,013	4,019	57%	2,736	39%	1	1	2	2	3	4	6	11	24	365	3	11.84	4		258	4%
924.00		<b>Thigh</b>	<b>2,352</b>	<b>1,412</b>	<b>60%</b>	<b>851</b>	<b>36%</b>	1	1	2	2	3	4	5	9	20	365	3	10.09	3		89	4%
924.01		<b>Hip</b>	<b>4,661</b>	<b>2,607</b>	<b>56%</b>	<b>1,885</b>	<b>40%</b>	1	1	2	3	3	4	6	12	26	365	3	12.62	4		169	4%
924.1	I	Knee and lower leg	23,852	15,220	64%	7,963	33%	1	1	2	2	3	4	5	8	19	365	3	9.27	3		669	3%
924.10		<b>Lower leg</b>	<b>7,089</b>	<b>4,553</b>	<b>64%</b>	<b>2,295</b>	<b>32%</b>	1	1	2	2	3	4	5	9	19	365	3	9.12	3		241	3%
924.11		<b>Knee</b>	<b>16,763</b>	<b>10,667</b>	<b>64%</b>	<b>5,668</b>	<b>34%</b>	1	1	2	2	3	4	5	8	19	365	3	9.33	3		428	3%
924.2	I	Ankle and foot, excluding toe(s)	13,722	8,615	63%	4,763	35%	1	1	2	2	3	4	5	7	16	365	3	7.89	3		344	3%
924.20		<b>Foot</b>	<b>10,226</b>	<b>6,529</b>	<b>64%</b>	<b>3,444</b>	<b>34%</b>	1	1	2	2	3	3	4	6	13	365	3	7.36	3		253	2%
924.21		<b>Ankle</b>	<b>3,496</b>	<b>2,086</b>	<b>60%</b>	<b>1,319</b>	<b>38%</b>	1	1	2	2	3	4	5	10	23	365	3	9.25	3		91	3%
924.3		<b>Toe</b>	<b>2,941</b>	<b>2,095</b>	<b>71%</b>	<b>779</b>	<b>26%</b>	1	1	1	2	2	3	4	5	10	365	2	5.58	3		67	2%
924.4		<b>Multiple sites of lower limb</b>	<b>69</b>	<b>40</b>	<b>58%</b>	<b>23</b>	<b>33%</b>	1	1	2	3	3	5	9	15	18	49	3	8.39	4		6	9%
926	I	Crushing injury of trunk	45	14	31%	23	51%	2	12	19	32	42	52	61	88	145	267	42	58.48	39		8	18%
926.1	I	Other specified sites	27	10	37%	16	59%	2	20	32	42	46	60	88	116	150	267	46	71.44	46		1	4%
927	I	Crushing injury of upper limb	6,769	4,572	68%	2,025	30%	1	1	2	3	4	6	11	19	37	365	4	15.70	6		172	3%
927.1	I	Elbow and forearm	205	98	48%	99	48%	1	2	3	4	6	10	14	26	86	364	6	27.12	6		8	4%

### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
927.10		Forearm	179	88	49%	83	46%	1	2	3	5	6	9	14	31	86	364	6	26.28	6		8	4%
927.11		Elbow	26	10	38%	16	62%	1	2	2	3	5	13	14	18	37	355	5	31.50	12			0%
927.2	I	Wrist and hand(s), except finger(s) alone	1,449	860	59%	545	38%	1	2	2	3	4	6	12	22	51	365	4	18.13	6		44	3%
927.20		Hand(s)	1,310	778	59%	491	37%	1	2	2	3	4	6	12	20	44	365	4	17.39	6		41	3%
927.21		Wrist	139	82	59%	54	39%	1	2	3	4	5	12	17	35	84	184	5	24.81	9		3	2%
927.3		Finger(s)	5,077	3,595	71%	1,362	27%	1	1	2	3	4	5	9	17	33	365	4	13.71	6		120	2%
928	I	Crushing injury of lower limb	1,350	584	43%	711	53%	1	2	3	5	9	13	24	37	74	365	9	27.71	18		55	4%
928.0	I	Hip and thigh	39	13	33%	25	64%	4	4	5	6	13	14	26	28	96	365	13	35.80	19		1	3%
928.00		Thigh	35	12	34%	22	63%	4	5	6	7	13	25	28	52	96	365	13	40.18	25		1	3%
928.1	I	Knee and lower leg	162	53	33%	102	63%	2	3	6	10	14	19	28	40	68	365	14	28.96	19		7	4%
928.10		Lower leg	131	38	29%	86	66%	2	3	6	10	13	19	28	37	68	365	13	29.21	20		7	5%
928.11		Knee	31	15	48%	16	52%	1	4	5	10	14	15	38	46	81	110	14	27.63	15			0%
928.2	I	Ankle and foot, excluding toe(s) alone	791	332	42%	422	53%	1	2	3	5	9	14	28	47	82	365	9	31.54	21		37	5%
928.20		Foot	708	304	43%	372	53%	1	2	3	4	8	13	25	47	81	365	8	31.27	20		32	5%
928.21		Ankle	83	28	34%	50	60%	3	6	8	10	13	28	34	44	95	174	13	33.52	28		5	6%
928.3		Toe(s)	353	185	52%	158	45%	1	2	3	4	5	7	13	20	34	126	5	13.38	13		10	3%
930	I	Foreign body on external eye	33,661	26,781	80%	5,973	18%	1	1	1	2	2	2	3	3	5	365	2	3.79	2		907	3%
930.0		Corneal foreign body	17,738	13,840	78%	3,390	19%	1	1	1	2	2	2	3	3	5	365	2	4.22	2		508	3%
930.1		Foreign body in conjunctival sac	3,116	2,675	86%	385	12%	1	1	1	1	2	2	2	3	5	63	2	2.97	2		56	2%
930.8		Other and combined sites	949	778	82%	146	15%	1	1	1	2	2	2	3	3	4	16	2	2.63	2		25	3%
930.9		Unspecified site	11,850	9,482	80%	2,050	17%	1	1	1	2	2	2	3	3	5	365	2	3.31	2		318	3%
931		Foreign body in ear	299	261	87%	26	9%	1	2	2	2	2	2	3	5	6	7	2	2.85	2		12	4%
940	I	Burn confined to eye and adnexa	3,073	2,035	66%	938	31%	1	1	1	2	2	2	3	4	6	365	2	4.23	2		100	3%
940.0		Chemical burn of eyelids and periocular area	341	235	69%	94	28%	1	1	1	2	2	2	3	4	5	66	2	3.78	2		12	4%
940.1		Other burns of eyelids and periocular area	290	215	74%	68	23%	1	1	1	1	1	2	2	3	4	37	1	2.51	2		7	2%
940.2		Alkaline chemical burn of cornea and conjunctival sac	238	162	68%	70	29%	1	2	2	3	4	4	6	11	18	179	4	11.09	4		6	3%
940.3		Acid chemical burn of cornea and conjunctival sac	215	148	69%	62	29%	1	1	1	2	2	3	3	4	6	48	2	3.61	2		5	2%
940.4		Other burn of cornea and conjunctival sac	845	538	64%	279	33%	1	1	1	2	2	3	3	4	6	212	2	3.80	2		28	3%
940.9		Unspecified burn of eye and adnexa	1,143	736	64%	365	32%	1	1	1	1	2	2	3	4	5	365	2	3.80	2		42	4%
941	I	Burn of face, head, and neck	3,191	2,016	63%	1,079	34%	1	1	2	3	4	5	8	13	23	365	4	10.84	5		96	3%
941.1	I	Erythema [first degree]	1,769	1,249	71%	467	26%	1	1	1	2	2	3	4	5	12	365	2	6.17	3		53	3%
941.10		Erythema [first degree]	727	491	68%	217	30%	1	1	1	2	2	3	3	5	10	365	2	5.96	3		19	3%
941.11		Erythema [first degree]	219	173	79%	41	19%	1	1	2	2	2	4	4	6	7	39	2	4.88	3		5	2%
941.12		Erythema [first degree]	80	50	63%	25	31%	1	1	1	2	2	2	3	4	5	8	2	2.48	3		5	6%
941.13		Erythema [first degree]	49	31	63%	16	33%	1	1	2	2	2	2	3	7	54	364	2	28.38	3		2	4%
941.14		Erythema [first degree]	43	23	53%	18	42%	1	2	3	4	4	8	13	14	26	33	4	9.00	3		2	5%
941.17		Erythema [first degree]	221	168	76%	46	21%	1	1	1	2	2	2	3	5	11	17	2	3.61	3		7	3%
941.18		Erythema [first degree]	277	204	74%	66	24%	1	1	1	2	2	3	4	5	13	98	2	5.77	3		7	3%
941.19		Erythema [first degree]	98	67	68%	28	29%	1	1	1	1	2	3	4	6	16	25	2	4.79	3		3	3%
941.2	I	Blisters, epidermal loss [second degree]	1,383	758	55%	587	42%	1	2	3	4	5	8	12	16	25	364	5	11.60	7		38	3%
941.20		Blisters, epidermal loss [second degree]	435	233	54%	190	44%	1	2	3	3	4	6	10	14	20	202	4	10.07	7		12	3%
941.21		Blisters, epidermal loss [second degree]	197	103	52%	87	44%	1	3	5	10	12	14	16	20	30	104	12	15.31	7		7	4%
941.22		Blisters, epidermal loss [second degree]	60	30	50%	28	47%	1	1	2	4	6	14	18	26	46	95	6	15.61	7		2	3%
941.25		Blisters, epidermal loss [second degree]	56	31	55%	25	45%	1	1	3	3	4	6	9	17	30	73	4	11.04	7			0%
941.27		Blisters, epidermal loss [second degree]	171	105	61%	63	37%	1	2	2	3	4	6	10	13	19	364	4	13.10	7		3	2%
941.28		Blisters, epidermal loss [second degree]	296	171	58%	118	40%	1	2	3	4	5	6	10	13	23	89	5	9.22	7		7	2%
941.29		Blisters, epidermal loss [second degree]	100	42	42%	53	53%	1	2	4	6	10	12	16	23	40	91	10	14.94	7		5	5%
941.3	I	Full-thickness skin loss [third degree NOS]	39	9	23%	25	64%	2	10	21	25	57	67	131	133	202	299	57	80.36	19		5	13%
942	I	Burn of trunk	1,382	816	59%	523	38%	1	1	2	3	4	6	12	22	38	365	4	16.28	5		43	3%
942.1	I	Erythema [first degree]	500	356	71%	130	26%	1	1	1	2	2	3	3	5	8	33	2	3.85	3		14	3%
942.12		Erythema [first degree]	182	131	72%	47	26%	1	1	1	2	3	3	4	5	11	20	3	4.09	3		4	2%

### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
942.13		Erythema [first degree]	136	94	69%	36	26%	1	1	2	2	2	3	3	4	6	33	2	3.78	3		6	4%
942.14		Erythema [first degree]	86	62	72%	20	23%	1	1	1	2	2	3	3	3	6	18	2	3.35	3		4	5%
942.2	I	Blisters, epidermal loss [second degree]	824	448	54%	351	43%	1	1	2	3	5	9	13	23	37	196	5	14.16	7		25	3%
942.20		Blisters, epidermal loss [second degree]	45	28	62%	17	38%	1	1	2	3	4	10	12	25	32	61	4	12.12	7			0%
942.21		Blisters, epidermal loss [second degree]	62	40	65%	21	34%	1	1	2	2	3	4	5	7	10	46	3	5.90	7		1	2%
942.22		Blisters, epidermal loss [second degree]	218	117	54%	95	44%	1	1	2	2	4	5	11	15	27	90	4	9.88	7		6	3%
942.23		Blisters, epidermal loss [second degree]	254	151	59%	95	37%	1	1	2	2	3	4	7	16	33	196	3	11.43	7		8	3%
942.24		Blisters, epidermal loss [second degree]	191	92	48%	92	48%	1	2	3	9	11	17	22	27	42	114	11	19.52	7		7	4%
942.25		Blisters, epidermal loss [second degree]	35	10	29%	24	69%	2	2	5	12	14	21	37	67	74	144	14	29.29	7		1	3%
942.3	I	Full-thickness skin loss [third degree NOS]	55	12	22%	39	71%	9	15	20	31	33	41	87	109	177	365	33	68.62	25		4	7%
942.34		Full-thickness skin loss [third degree NOS]	19	2	11%	16	84%	14	17	18	31	33	38	40	79	169	365	33	62.63	25		1	5%
943	I	Burn of upper limb, except wrist and hand	5,683	3,904	69%	1,627	29%	1	1	2	2	3	4	5	10	20	365	3	10.22	3		152	3%
943.1	I	Erythema [first degree]	1,936	1,476	76%	416	21%	1	1	1	2	2	2	3	4	8	365	2	5.81	3		44	2%
943.10		Erythema [first degree]	260	195	75%	57	22%	1	1	1	2	2	3	4	6	8	365	2	9.96	3		8	3%
943.11		Erythema [first degree]	1,396	1,066	76%	299	21%	1	1	1	2	2	2	3	4	6	364	2	5.02	3		31	2%
943.13		Erythema [first degree]	120	95	79%	25	21%	1	1	1	2	2	2	2	4	33	41	2	6.28	3			0%
943.15		Erythema [first degree]	72	53	74%	17	24%	1	1	1	1	1	2	2	5	18	20	1	4.18	3		2	3%
943.2	I	Blisters, epidermal loss [second degree]	3,600	2,377	66%	1,125	31%	1	1	2	2	3	4	6	10	18	365	3	8.10	4		98	3%
943.20		Blisters, epidermal loss [second degree]	340	221	65%	101	30%	1	1	1	2	3	4	6	8	16	24	3	5.23	4		18	5%
943.21		Blisters, epidermal loss [second degree]	2,679	1,810	68%	808	30%	1	1	2	2	3	4	5	9	16	365	3	7.36	4		61	2%
943.22		Blisters, epidermal loss [second degree]	154	108	70%	41	27%	1	1	2	2	3	4	6	14	34	163	3	13.78	4		5	3%
943.23		Blisters, epidermal loss [second degree]	272	167	61%	97	36%	1	1	2	2	3	5	10	18	26	131	3	11.70	4		8	3%
943.25		Blisters, epidermal loss [second degree]	87	44	51%	40	46%	1	1	1	1	2	3	3	6	12	42	2	5.58	4		3	3%
943.29		Blisters, epidermal loss [second degree]	43	18	42%	24	56%	1	3	4	8	13	20	22	27	39	131	13	20.04	4		1	2%
943.3	I	Full-thickness skin loss [third degree NOS]	143	50	35%	83	58%	2	7	14	23	30	39	48	74	120	365	30	56.52	23		10	7%
943.31		Full-thickness skin loss [third degree NOS]	87	32	37%	47	54%	1	2	9	14	22	30	41	58	90	270	22	40.13	23		8	9%
944	I	Burn of wrist(s) and hand(s)	9,346	6,035	65%	3,062	33%	1	1	2	2	3	4	5	8	17	365	3	8.02	3		249	3%
944.1	I	Erythema [first degree]	3,076	2,204	72%	804	26%	1	1	1	2	2	2	3	4	6	364	2	4.35	2	2	68	2%
944.10		Erythema [first degree]	1,705	1,190	70%	481	28%	1	1	1	2	2	3	3	4	6	364	2	4.48	2		34	2%
944.11		Erythema [first degree]	436	341	78%	86	20%	1	1	1	1	2	2	3	4	6	30	2	3.53	2		9	2%
944.12		Erythema [first degree]	112	88	79%	21	19%	1	1	1	1	1	2	3	3	4	5	1	2.00	2		3	3%
944.13		Erythema [first degree]	89	65	73%	22	25%	1	1	1	2	2	3	3	4	6	364	2	19.14	2		2	2%
944.15		Erythema [first degree]	185	128	69%	54	29%	1	1	1	2	2	3	3	5	6	13	2	3.04	2		3	2%
944.16		Erythema [first degree]	167	107	64%	57	34%	1	1	1	2	2	3	3	5	6	8	2	2.70	2		3	2%
944.17		Erythema [first degree]	276	210	76%	58	21%	1	1	1	1	2	2	2	4	6	13	2	2.48	2		8	3%
944.18		Erythema [first degree]	61	41	67%	15	25%	1	2	2	2	3	3	4	5	17	37	3	6.07	2		5	8%
944.2	I	Blisters, epidermal loss [second degree]	6,037	3,743	62%	2,122	35%	1	1	2	2	3	4	5	9	16	365	3	6.94	4	5	172	3%
944.20		Blisters, epidermal loss [second degree]	2,571	1,569	61%	917	36%	1	1	2	2	3	3	4	6	12	174	3	5.28	4		85	3%
944.21		Blisters, epidermal loss [second degree]	947	662	70%	264	28%	1	1	2	2	3	4	5	9	16	107	3	6.36	4		21	2%
944.22		Blisters, epidermal loss [second degree]	272	185	68%	83	31%	1	1	2	2	3	4	5	7	14	365	3	9.39	4		4	1%
944.23		Blisters, epidermal loss [second degree]	324	191	59%	123	38%	1	1	1	2	2	4	6	9	16	110	2	6.96	4		10	3%
944.24		Blisters, epidermal loss [second degree]	175	111	63%	63	36%	1	1	2	3	4	5	10	19	38	365	4	16.71	4		1	1%
944.25		Blisters, epidermal loss [second degree]	436	239	55%	186	43%	1	1	2	3	4	6	7	12	20	42	4	7.60	4		11	3%
944.26		Blisters, epidermal loss [second degree]	502	271	54%	217	43%	1	1	2	3	5	8	11	16	22	86	5	9.76	4		14	3%
944.27		Blisters, epidermal loss [second degree]	623	415	67%	188	30%	1	1	2	2	2	3	4	7	17	140	2	6.69	4		20	3%
944.28		Blisters, epidermal loss [second degree]	187	100	53%	81	43%	1	2	3	3	4	8	11	14	18	61	4	9.00	4		6	3%
944.3	I	Full-thickness skin loss [third degree NOS]	227	86	38%	134	59%	2	4	10	18	25	33	44	59	106	365	25	42.69	20	7	7	3%
944.30		Full-thickness skin loss [third degree NOS]	72	29	40%	39	54%	1	3	5	11	23	33	44	83	171	316	23	50.15	20		4	6%
944.31		Full-thickness skin loss [third degree NOS]	59	29	49%	28	47%	2	4	9	21	25	34	48	54	80	112	25	33.14	20		2	3%
944.36		Full-thickness skin loss [third degree NOS]	25	5	20%	20	80%	5	15	18	31	33	38	39	55	153	365	33	59.75	20			0%
945	I	Burn of lower limb(s)	3,502	1,866	53%	1,537	44%	1	2	3	4	6	12	19	30	52	365	6	19.91	14		99	3%



### Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
945.1	I	Erythema [first degree]	881	626	71%	227	26%	1	1	2	2	2	3	4	6	12	58	2	5.25	4		28	3%
945.10		<b>Erythema [first degree]</b>	122	91	75%	29	24%	1	1	1	1	3	3	4	5	23	58	3	6.79	4		2	2%
945.12		<b>Erythema [first degree]</b>	302	207	69%	82	27%	1	1	2	2	3	3	5	7	13	36	3	5.41	4		13	4%
945.14		<b>Erythema [first degree]</b>	163	121	74%	38	23%	1	1	1	1	2	2	3	5	10	21	2	3.84	4		4	2%
945.15		<b>Erythema [first degree]</b>	59	38	64%	20	34%	1	1	2	2	2	3	3	6	9	36	2	5.05	4		1	2%
945.16		<b>Erythema [first degree]</b>	139	102	73%	34	24%	1	1	2	2	2	2	2	4	5	33	2	3.68	4		3	2%
945.2	I	Blisters, epidermal loss [second degree]	2,312	1,154	50%	1,102	48%	1	2	3	4	6	11	17	25	40	365	6	16.80	9		56	2%
945.20		<b>Blisters, epidermal loss [second degree]</b>	217	131	60%	75	35%	1	2	2	3	5	9	14	28	47	365	5	22.87	9		11	5%
945.21		<b>Blisters, epidermal loss [second degree]</b>	60	29	48%	30	50%	1	1	2	2	4	13	28	35	58	141	4	22.83	9		1	2%
945.22		<b>Blisters, epidermal loss [second degree]</b>	769	364	47%	384	50%	1	2	3	4	6	8	14	19	30	227	6	13.33	9		21	3%
945.23		<b>Blisters, epidermal loss [second degree]</b>	406	175	43%	224	55%	1	2	4	5	10	17	24	34	49	143	10	18.88	9		7	2%
945.24		<b>Blisters, epidermal loss [second degree]</b>	467	259	55%	198	42%	1	2	3	4	6	12	17	24	36	144	6	15.54	9		10	2%
945.25		<b>Blisters, epidermal loss [second degree]</b>	77	47	61%	29	38%	1	2	2	4	5	6	7	10	18	121	5	9.59	9		1	1%
945.26		<b>Blisters, epidermal loss [second degree]</b>	282	141	50%	137	49%	1	2	3	4	9	11	16	27	44	164	9	18.66	9		4	1%
945.29		<b>Blisters, epidermal loss [second degree]</b>	34	8	24%	25	74%	2	4	6	13	15	40	60	67	83	89	15	34.28	9		1	3%
945.3	I	Full-thickness skin loss [third degree NOS]	299	85	28%	199	67%	4	11	21	29	34	44	56	74	102	365	34	48.96	30		15	5%
945.32		<b>Full-thickness skin loss [third degree NOS]</b>	113	31	27%	77	68%	5	12	18	26	33	40	45	62	82	131	33	38.65	30		5	4%
945.33		<b>Full-thickness skin loss [third degree NOS]</b>	64	19	30%	43	67%	3	13	26	38	40	47	58	72	89	175	40	45.98	30		2	3%
945.34		<b>Full-thickness skin loss [third degree NOS]</b>	60	17	28%	40	67%	4	16	24	29	41	52	58	75	159	355	41	65.70	30		3	5%
948	I	Burns classified according to extent of body surface involved	236	137	58%	86	36%	1	1	1	2	3	5	9	40	231	365	3	50.41	6	1	13	6%
948.0	I	Burn [any degree] involving less than 10 percent of body surface	205	130	63%	66	32%	1	1	1	2	2	2	4	5	14	205	2	7.33	5		9	4%
948.00		<b>Burn [any degree] involving less than 10 percent of body surface</b>	197	127	64%	61	31%	1	1	1	2	2	2	4	5	9	40	2	4.48	5		9	5%
953	I	Injury to nerve roots and spinal plexus	57	6	11%	38	67%	4	10	19	57	67	112	178	270	365	365	67	123.55	26		13	23%
953.4		<b>Brachial plexus</b>	37	2	5%	28	76%	4	11	20	57	66	112	157	296	365	365	66	125.89	57		7	19%
955	I	Injury to peripheral nerve(s) of shoulder girdle and upper limb	852	201	24%	592	69%	5	12	19	26	37	54	70	95	178	365	37	68.52	33		59	7%
955.1		<b>Median nerve</b>	76	16	21%	54	71%	11	17	31	53	67	87	114	193	365	365	67	109.98	53		6	8%
955.2		<b>Ulnar nerve</b>	181	62	34%	99	55%	3	7	18	26	39	62	114	157	298	365	39	88.33	32		20	11%
955.3		<b>Radial nerve</b>	96	16	17%	75	78%	3	13	20	29	38	54	72	89	161	365	38	66.60	35		5	5%
955.6		<b>Digital nerve</b>	452	103	23%	330	73%	5	12	17	24	31	45	60	78	104	365	31	53.02	33		19	4%
956	I	Injury to peripheral nerve(s) of pelvic girdle and lower limb	57	12	21%	36	63%	16	25	31	45	59	75	131	184	222	365	59	95.31	9		9	16%
956.3		<b>Peroneal nerve</b>	26	4	15%	18	69%	4	18	34	59	65	84	91	137	222	226	65	86.50	9		4	15%
957	I	Injury to other and unspecified nerves	30	12	40%	16	53%	6	12	19	39	40	48	102	125	238	365	40	80.25	14	18	2	7%
957.1		<b>Other specified nerve(s)</b>	27	10	37%	15	56%	6	12	37	39	44	48	102	125	238	365	44	84.87	40		2	7%
958	I	Certain early complications of trauma	261	110	42%	134	51%	2	7	12	30	57	79	106	174	278	365	57	93.60	13		17	7%
958.3		<b>Posttraumatic wound infection, not elsewhere classified</b>	147	101	69%	44	30%	1	1	2	5	7	9	12	24	52	178	7	17.93	6		2	1%
958.8		<b>Other early complications of trauma</b>	100	5	5%	83	83%	11	37	57	75	93	116	174	226	365	365	93	131.49	54		12	12%
982	I	Toxic effect of solvents other than petroleum based	75	54	72%	18	24%	1	1	2	2	2	3	3	4	8	2	2.67	2			3	4%
983	I	Toxic effect of corrosive aromatics, acids, and caustic alkalis	170	123	72%	41	24%	1	1	1	2	2	3	3	4	6	12	2	2.93	2		6	4%
983.9		<b>Caustic, unspecified</b>	73	48	66%	23	32%	1	1	1	2	2	2	3	5	6	12	2	2.87	2		2	3%
984	I	Toxic effect of lead and its compounds (including fumes)	33	15	45%	15	45%	7	19	35	39	58	59	77	123	216	244	58	77.27	2		3	9%
986		<b>Toxic effect of carbon monoxide</b>	461	303	66%	140	30%	1	1	1	1	2	2	3	4	5	62	2	3.54	2		18	4%
987	I	Toxic effect of other gases, fumes, or vapors	3,240	2,378	73%	744	23%	1	1	1	2	2	3	3	5	8	365	2	7.55	2		118	4%
987.6		<b>Chlorine gas</b>	156	112	72%	39	25%	1	1	1	2	2	2	4	5	6	365	2	12.69	2		5	3%
987.8		<b>Other specified gases, fumes, or vapors</b>	503	377	75%	103	20%	1	1	2	2	3	3	4	5	10	60	3	4.96	3		23	5%
987.9		<b>Unspecified gas, fume, or vapor</b>	2,458	1,801	73%	569	23%	1	1	1	2	2	3	3	5	7	365	2	7.07	2		88	4%
989	I	Toxic effect of other substances, chiefly nonmedicinal as to source	5,205	4,259	82%	803	15%	1	1	1	1	2	2	3	3	5	365	2	4.39	2		143	3%
989.5		<b>Venom</b>	4,932	4,064	82%	738	15%	1	1	1	1	2	2	2	3	5	365	2	3.22	2	2	130	3%
989.8	I	Other substances, chiefly nonmedicinal as to source	156	112	72%	35	22%	1	1	1	2	2	3	4	6	72	364	2	28.46	2	3	9	6%
989.89		<b>Other</b>	127	96	76%	26	20%	1	1	1	1	2	3	4	5	31	364	2	21.27	2		5	4%
991	I	Effects of reduced temperature	150	87	58%	49	33%	1	1	2	3	3	4	7	12	41	74	3	10.82	4	5	14	9%
991.1		<b>Frostbite of hand</b>	67	35	52%	25	37%	1	1	2	2	3	5	13	23	57	74	3	14.88	5		7	10%

**Ohio Specific Workers' Compensation Disability Duration Outcomes per ICD-9 Code**

ICD9 Code	Invalid ICD-9	Name	No. of Claims	0-day Cases	% 0 days	Lost-day Cases	% Lost-day Cases	10th %	20th %	30th %	40th %	50th %	60th %	70th %	80th %	90th %	100th %	Ohio Med.	Ohio Mean	ODG Med.	OSHA Med.	Open Cases	% Open
992	I	Effects of heat and light	895	593	66%	283	32%	1	1	1	1	2	2	3	4	5	56	2	3.30	2	2	19	2%
<b>992.5</b>		<b>Heat exhaustion, unspecified</b>	<b>823</b>	<b>544</b>	<b>66%</b>	<b>263</b>	<b>32%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>31</b>	<b>2</b>	<b>2.81</b>	<b>3</b>		<b>16</b>	<b>2%</b>
993	I	Effects of air pressure	104	77	74%	24	23%	1	1	2	2	3	4	8	15	16	21	3	6.54	3	7	3	3%
994	I	Effects of other external causes	2,365	1,685	71%	588	25%	1	1	1	2	2	3	3	5	11	365	2	8.41	2		92	4%
<b>994.8</b>		<b>Electrocution and nonfatal effects of electric current</b>	<b>1,303</b>	<b>880</b>	<b>68%</b>	<b>373</b>	<b>29%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>11</b>	<b>365</b>	<b>2</b>	<b>7.51</b>	<b>3</b>	<b>4</b>	<b>50</b>	<b>4%</b>
<b>994.9</b>		<b>Other effects of external causes</b>	<b>1,008</b>	<b>772</b>	<b>77%</b>	<b>198</b>	<b>20%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>7</b>	<b>365</b>	<b>2</b>	<b>8.33</b>	<b>2</b>		<b>38</b>	<b>4%</b>
995	I	Certain adverse effects not elsewhere classified	366	256	70%	92	25%	1	1	1	2	2	3	4	5	7	354	2	9.01	4		18	5%
<b>995.2</b>		<b>Unspecified adverse effect of drug, medicinal and biological substance</b>	<b>33</b>	<b>15</b>	<b>45%</b>	<b>17</b>	<b>52%</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>5</b>	<b>5</b>	<b>7</b>	<b>14</b>	<b>16</b>	<b>3</b>	<b>4.71</b>	<b>13</b>		<b>1</b>	<b>3%</b>
<b>995.3</b>		<b>Allergy, unspecified</b>	<b>296</b>	<b>221</b>	<b>75%</b>	<b>61</b>	<b>21%</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>156</b>	<b>2</b>	<b>5.20</b>	<b>3</b>	<b>7</b>	<b>14</b>	<b>5%</b>
996	I	Complications peculiar to certain specified procedures	212	17	8%	133	63%	20	44	79	109	143	181	302	365	365	365	143	180.50	10		62	29%
996.4	I	Mechanical complication of internal orthopedic device, implant, and graft	69	4	6%	43	62%	19	34	54	92	109	167	278	365	365	365	109	166.14	91	167	22	32%
996.6	I	Infection and inflammatory reaction due to internal prosthetic device, implan	23		0%	16	70%	31	44	71	99	135	175	216	224	365	365	135	158.63	15		7	30%
996.7	I	Other complications of internal (biological) (synthetic) prosthetic device, imp	89	3	3%	58	65%	44	89	123	146	181	248	365	365	365	365	181	213.97	15	30	28	31%
<b>996.78</b>		<b>Due to other internal orthopedic device, implant, and graft</b>	<b>74</b>	<b>3</b>	<b>4%</b>	<b>50</b>	<b>68%</b>	<b>27</b>	<b>88</b>	<b>111</b>	<b>141</b>	<b>166</b>	<b>205</b>	<b>322</b>	<b>365</b>	<b>365</b>	<b>365</b>	<b>166</b>	<b>201.46</b>	<b>90</b>		<b>21</b>	<b>28%</b>
997	I	Complications affecting specified body systems, not elsewhere classified	21		0%	17	81%	1	5	31	35	53	90	109	335	365	365	53	124.24	11		4	19%
998	I	Other complications of procedures, NEC	191	52	27%	100	52%	10	27	45	70	91	140	194	313	365	365	91	142.95	13		39	20%
998.5	I	Postoperative infection	101	9	9%	59	58%	41	52	81	107	161	194	313	358	365	365	161	181.81	15		33	33%
<b>998.59</b>		<b>Other postoperative infection</b>	<b>94</b>	<b>8</b>	<b>9%</b>	<b>56</b>	<b>60%</b>	<b>41</b>	<b>52</b>	<b>78</b>	<b>106</b>	<b>143</b>	<b>194</b>	<b>313</b>	<b>358</b>	<b>365</b>	<b>365</b>	<b>143</b>	<b>178.95</b>	<b>67</b>		<b>30</b>	<b>32%</b>